

Self-interacting diffusions IV: Rate of convergence*

Michel Benaïm

Université de Neuchâtel, Suisse

Olivier Raimond

Université Paris Ouest Nanterre la défense, France

July 31, 2009

Abstract

Self-interacting diffusions are processes living on a compact Riemannian manifold defined by a stochastic differential equation with a drift term depending on the past empirical measure μ_t of the process. The asymptotics of μ_t is governed by a deterministic dynamical system and under certain conditions (μ_t) converges almost surely towards a deterministic measure μ^* (see Benaïm, Ledoux, Raimond (2002) and Benaïm, Raimond (2005)). We are interested here in the rate of convergence of μ_t towards μ^* . A central limit theorem is proved. In particular, this shows that greater is the interaction repelling faster is the convergence.

*We acknowledge financial support from the Swiss National Science Foundation Grant 200021-103625/1

1 Introduction

Self-interacting diffusions

Let M be a smooth compact Riemannian manifold and $V : M \times M \rightarrow \mathbb{R}$ a sufficiently smooth mapping¹. For all finite Borel measure μ , let $V\mu : M \rightarrow \mathbb{R}$ be the smooth function defined by

$$V\mu(x) = \int_M V(x, y)\mu(dy).$$

Let (e_α) be a finite family of vector fields on M such that

$$\sum_{\alpha} e_{\alpha}(e_{\alpha}f)(x) = \Delta f(x),$$

where Δ is the Laplace operator on M and $e_{\alpha}(f)$ stands for the Lie derivative of f along e_{α} . Let (B^{α}) be a family of independent Brownian motions.

A *self-interacting diffusion* on M associated to V can be defined as the solution to the stochastic differential equation (SDE)

$$dX_t = \sum_{\alpha} e_{\alpha}(X_t) \circ dB_t^{\alpha} - \nabla(V\mu_t)(X_t)dt.$$

where

$$\mu_t = \frac{1}{t} \int_0^t \delta_{X_s} ds$$

is the empirical occupation measure of (X_t) .

In absence of drift (i.e $V = 0$), (X_t) is just a Brownian motion on M but in general it defines a non Markovian process whose behavior at time t depends on its past trajectories through μ_t . This type of process was introduced in Benaim, Ledoux and Raimond (2002) (hence after referred as [3]) and further analyzed in a series of papers by Benaim and Raimond (2003, 2005, 2007) (hence after referred as [4], [5] and [6]). We refer the reader to these papers for more details and especially to [3] for a detailed construction of the process and its elementary properties. For a general overview of processes with reinforcement we refer the reader to the recent survey paper by Pemantle (2007) ([15]).

¹The mapping $V_x : M \rightarrow \mathbb{R}$ defined by $V_x(y) = V(x, y)$ is C^2 and its derivatives are continuous in (x, y)

Notation and Background

Standing Notation We let $\mathcal{M}(M)$ denote the space of finite Borel measures on M , $\mathcal{P}(M) \subset \mathcal{M}(M)$ the space of probability measures. If I is a metric space (typically, $I = M, \mathbb{R}^+ \times M$ or $[0, T] \times M$) we let $C(I)$ denote the space of real valued continuous functions on I equipped with the topology of uniform convergence on compact sets. When I is compact and $f \in C(I)$ we let $\|f\| = \sup_{x \in I} |f(x)|$. The normalized Riemann measure on M will be denoted by λ .

Let $\mu \in \mathcal{P}(M)$ and $f : M \rightarrow \mathbb{R}$ a nonnegative or μ -integrable Borel function. We write μf for $\int f d\mu$, and $f\mu$ for the measure defined as $f\mu(A) = \int_A f d\mu$. We let $L^2(\mu)$ denote the space of such functions for which $\mu|f|^2 < \infty$, equipped with the inner product

$$\langle f, g \rangle_\mu = \mu(fg)$$

and the norm

$$\|f\|_\mu = \sqrt{\mu f^2}.$$

We simply write L^2 for $L^2(\lambda)$.

Of fundamental importance in the analysis of the asymptotics of (μ_t) is the mapping $\Pi : \mathcal{M}(M) \rightarrow \mathcal{P}(M)$ defined by

$$\Pi(\mu) = \xi(V\mu)\lambda \tag{1}$$

where $\xi : C(M) \rightarrow C(M)$ is the function defined by

$$\xi(f)(x) = \frac{e^{-f(x)}}{\int_M e^{-f(y)} \lambda(dy)}. \tag{2}$$

In [3], it is shown that the asymptotics of μ_t can be precisely related to the long term behavior of a certain semiflow on $\mathcal{P}(M)$ induced by the ordinary differential equation (ODE) on $\mathcal{M}(M)$:

$$\dot{\mu} = -\mu + \Pi(\mu). \tag{3}$$

Depending on the nature of V , the dynamics of (3) can either be convergent or nonconvergent leading to similar behaviors for $\{\mu_t\}$ (see [3]). When V is symmetric, (3) happens to be a *quasigradient* and the following convergence result hold.

Theorem 1.1 ([5]) *Assume that V is symmetric. i.e. $V(x, y) = V(y, x)$. Then the limit set of $\{\mu_t\}$ (for the topology of weak* convergence) is almost surely a compact connected subset of*

$$\text{Fix}(\Pi) = \{\mu \in \mathcal{P}(M) : \mu = \Pi(\mu)\}.$$

In particular, if $\text{Fix}(\Pi)$ is finite then (μ_t) converges almost surely toward a fixed point of Π . This holds for a generic function V (see [5]).

Sufficient conditions ensuring that $\text{Fix}(\Pi)$ has cardinal one are as follows:

Theorem 1.2 ([5], [6]) *Assume that V is symmetric and that one of the two following conditions hold*

(i) *Up to an additive constant V is a Mercer kernel, That is*

$$V(x, y) = K(x, y) + C$$

and

$$\int K(x, y) f(x) f(y) \lambda(dx) \lambda(dy) \geq 0$$

for all $f \in L^2$.

(ii) *For all $x \in M, y \in M, u \in T_x M, v \in T_y M$*

$$\text{Ric}_x(u, u) + \text{Ric}_y(v, v) + \text{Hess}_{x,y} V((u, v), (u, v)) \geq K(\|u\|^2 + \|v\|^2)$$

where K is some positive constant. Here Ric_x stands for the Ricci tensor at x and $\text{Hess}_{x,y}$ is the Hessian of V at (x, y) .

Then $\text{Fix}(\Pi)$ reduces to a singleton $\{\mu^\}$ and $\mu_t \rightarrow \mu^*$ with probability one.*

As observed in [6] the condition (i) in Theorem 1.2 seems well suited to describe *self-repelling diffusions*. On the other hand, it is not clearly related to the geometry of M . Condition (ii) has a more geometrical flavor and is robust to smooth perturbations (of M and V). It can be seen as a Bakry-Emery type condition for self interacting diffusions.

In [5], it is also proved that every stable (for the ODE (3)) fixed point of Π has a positive probability to be a limit point for μ_t ; and any unstable fixed point cannot be a limit point for μ_t .

Organisation of the paper

Let $\mu^* \in \text{Fix}(\Pi)$. We will assume that

Hypothesis 1.3 μ_t converges a.s. towards μ^* .

Sufficient conditions are given by Theorem 1.2

In this paper we intend to study the rate of this convergence. Let

$$\Delta_t = e^{t/2}(\mu_{e^t} - \mu^*).$$

It will be shown that, under some conditions to be specified later, for all $g = (g_1, \dots, g_n) \in C(M)^n$ the process

$$[\Delta_s g_1, \dots, \Delta_s g_n, V \Delta_s]_{s \geq t}$$

converges in law, as $t \rightarrow \infty$, toward a certain stationary Ornstein-Uhlenbeck process (Z^g, Z) on $\mathbb{R}^n \times C(M)$. This process is defined in Section 2. The main result is stated in section 3 and some examples are developed. It is in particular observed that a strong repelling interaction gives a faster convergence. The section 4 is a proof section. The appendix, section 5, contains general material on random variables and Ornstein-Uhlenbeck processes on $C(M)$.

In the following K (respectively C) denotes a positive constant (respectively a positive random constant). These constants may change from line to line.

2 The Ornstein-Uhlenbeck process (Z^g, Z) .

Throughout all this section we let $\mu \in \mathcal{P}(M)$. For $x \in M$ we set $V_x : M \rightarrow \mathbb{R}$ defined by $V_x(y) = V(x, y)$.

2.1 The operator G_μ

Let $g \in C(M)$ and let $G_{\mu, g} : \mathbb{R} \times C(M) \rightarrow \mathbb{R}$ be the linear operator defined by

$$G_{\mu, g}(u, f) = u/2 + \text{Cov}_\mu(g, f), \quad (4)$$

where Cov_μ is the covariance on $L^2(\mu)$, that is the bilinear form acting on $L^2 \times L^2$ defined by

$$\text{Cov}_\mu(f, g) = \mu(fg) - (\mu f)(\mu g).$$

We define the linear operator $G_\mu : C(M) \rightarrow C(M)$ by

$$\begin{aligned} G_\mu f(x) &= G_{\mu, V_x}(f(x), f) \\ &= f(x)/2 + \text{Cov}_\mu(V_x, f). \end{aligned} \tag{5}$$

It is easily seen that $\|G_\mu f\| \leq (2\|V\| + 1/2)\|f\|$. In particular, G_μ is a bounded operator. Let $\{e^{-tG_\mu}\}$ denotes the semigroup acting on $C(M)$ with generator $-G_\mu$. From now on we will assume the following:

Hypothesis 2.1 *There exists $\kappa > 0$ and $\hat{\lambda} \in \mathcal{P}(M)$ such that $\mu < \hat{\lambda}$ with $\|\frac{d\mu}{d\hat{\lambda}}\|_\infty < \infty$, λ and $\hat{\lambda}$ are equivalent measures with $\|\frac{d\lambda}{d\hat{\lambda}}\|_\infty < \infty$ and $\|\frac{d\hat{\lambda}}{d\lambda}\|_\infty < \infty$, and such that for all $f \in L^2(\hat{\lambda})$,*

$$\langle G_\mu f, f \rangle_{\hat{\lambda}} \geq \kappa \|f\|_{\hat{\lambda}}^2.$$

Let

$$\lambda(-G_\mu) = \lim_{t \rightarrow \infty} \frac{\log(\|e^{-tG_\mu}\|)}{t}.$$

This limit exists by subadditivity. Then

Lemma 2.2 *Hypothesis 2.1 implies that $\lambda(-G_\mu) \leq -\kappa < 0$.*

Proof : For all $f \in L^2(\hat{\lambda})$,

$$\begin{aligned} \frac{d}{dt} \|e^{-tG_\mu} f\|_{\hat{\lambda}}^2 &= -2 \langle G_\mu e^{-tG_\mu} f, e^{-tG_\mu} f \rangle_{\hat{\lambda}} \\ &\leq -2\kappa \|e^{-tG_\mu} f\|_{\hat{\lambda}}^2. \end{aligned}$$

This implies that $\|e^{-tG_\mu} f\|_{\hat{\lambda}} \leq e^{-\kappa t} \|f\|_{\hat{\lambda}}$.

Denote by g_t the solution of the differential equation

$$\frac{dg_t}{dt} = \text{Cov}_\mu(V_x, g_t)$$

with $g_0 = f$, where $f \in C(M)$. Note that $e^{-tG_\mu} f = e^{-t/2} g_t$. It is straightforward to check that (using the fact that $\|\frac{d\mu}{d\hat{\lambda}}\|_\infty < \infty$)

$$\frac{d}{dt} \|g_t\|_{\hat{\lambda}} \leq K \|g_t\|_{\hat{\lambda}}$$

with K a constant depending only on V and μ . Thus

$$\sup_{t \in [0,1]} \|g_t\|_{\hat{\lambda}} \leq K \|f\|_{\hat{\lambda}}.$$

Now, since for all $x \in M$ and $t \in [0, 1]$

$$\left| \frac{d}{dt} g_t(x) \right| \leq K \|g_t\|_{\hat{\lambda}} \leq K \|f\|_{\hat{\lambda}},$$

we have $\|g_1\| \leq K \|f\|_{\hat{\lambda}}$. This implies that

$$\|e^{-G_\mu} f\| \leq K \|f\|_{\hat{\lambda}}.$$

Now for all $t > 1$, and $f \in C(M)$,

$$\begin{aligned} \|e^{-tG_\mu} f\| &= \|e^{-G_\mu} e^{-(t-1)G_\mu} f\| \\ &\leq K \|e^{-(t-1)G_\mu} f\|_{\hat{\lambda}} \\ &\leq K e^{-\kappa(t-1)} \|f\|_{\hat{\lambda}} \\ &\leq K e^{-\kappa t} \|f\|_{\infty}. \end{aligned}$$

This implies that $\|e^{-tG_\mu}\| \leq K e^{-\kappa t}$, which proves the lemma. **QED**

The *adjoint* of G_μ is the operator on $\mathcal{M}(M)$ defined by the relation

$$m(G_\mu f) = (G_\mu^* m) f$$

for all $m \in \mathcal{M}(M)$ and $f \in C(M)$. It is not hard to verify that

$$G_\mu^* m = \frac{1}{2} m + (Vm)\mu - (\mu(Vm))\mu. \quad (6)$$

2.2 The generator A_μ and its inverse Q_μ

Let H^2 be the Sobolev space of real valued functions on M , associated with the norm $\|f\|_H^2 = \|f\|_\lambda^2 + \|\nabla f\|_\lambda^2$. Since $\Pi(\mu)$ and λ are equivalent measures with continuous Radon-Nykodim derivative, $L^2(\Pi(\mu)) = L^2(\lambda) := L^2$. We denote by K_μ the projection operator, acting on $L^2(\Pi(\mu))$, defined by

$$K_\mu f = f - \Pi(\mu) f.$$

We denote by A_μ the operator acting on H^2 defined by

$$A_\mu f = \frac{1}{2} \Delta f - \langle \nabla V \mu, \nabla f \rangle.$$

Note that for f and g in L^2 ,

$$\langle A_\mu f, g \rangle_{\Pi(\mu)} = -\frac{1}{2} \int \langle \nabla f, \nabla g \rangle(x) \Pi(\mu)(dx)$$

where $\langle \cdot, \cdot \rangle$ denotes the Riemannian inner product on M .

For all $f \in C(M)$ there exists $\mathbf{Q}_\mu f \in H^2$ such that $\Pi(\mu)(\mathbf{Q}_\mu f) = 0$ and

$$f - \Pi(\mu)f = K_\mu f = -A_\mu \mathbf{Q}_\mu f. \quad (7)$$

Note that if P_t^μ denotes the semigroup with generator A_μ , then

$$\mathbf{Q}_\mu f = \int_0^\infty P_t^\mu K_\mu f dt.$$

Since there exists $p_t^\mu(\cdot, \cdot)$ such that

$$P_t^\mu f(x) = \int_M p_t^\mu(x, y) f(y) \Pi(\mu)(dy),$$

we have

$$\mathbf{Q}_\mu f(x) = \int_M q_\mu(x, y) f(y) \Pi(\mu)(dy)$$

where

$$q_\mu(x, y) = \int_0^\infty (p_t^\mu(x, y) - 1) dt.$$

Then, as shown in [3], $\mathbf{Q}_\mu f$ is C^1 and there exists a constant K such that for all $f \in C(M)$ and $\mu \in \mathcal{P}(M)$,

$$\|\mathbf{Q}_\mu f\|_\infty \leq K \|f\|_\infty \quad (8)$$

$$\|\nabla \mathbf{Q}_\mu f\|_\infty \leq K \|f\|_\infty. \quad (9)$$

Finally, note that for f and g in L^2 ,

$$\begin{aligned} \int \langle \nabla \mathbf{Q}_\mu f, \nabla \mathbf{Q}_\mu g \rangle(x) \Pi(\mu)(dx) &= -2 \langle A_\mu \mathbf{Q}_\mu f, \mathbf{Q}_\mu g \rangle_{\Pi(\mu)} \\ &= 2 \langle f, \mathbf{Q}_\mu g \rangle_{\Pi(\mu)}. \end{aligned} \quad (10)$$

2.3 The covariance C_μ

We let $\widehat{C}_\mu$ denote the bilinear continuous form $\widehat{C}_\mu : C(M) \times C(M) \rightarrow \mathbb{R}$ defined by

$$\widehat{C}_\mu(f, g) = 2\langle f, \mathbf{Q}_\mu g \rangle_{\Pi(\mu)}.$$

This form is symmetric (see its expression given by (10)). Note also that for some constant depending on μ ,

$$|\widehat{C}_\mu(f, g)| \leq K\|f\| \times \|g\|.$$

We let C_μ denote the mapping $C_\mu : M \times M \rightarrow \mathbb{R}$ defined by

$$C_\mu(x, y) = \widehat{C}_\mu(V_x, V_y).$$

Then C_μ is a covariance function (or a Mercer kernel), i.e. it is continuous, symmetric and $\sum_{i,j} \lambda_i \lambda_j C_\mu(x_i, x_j) \geq 0$.

2.4 The process Z

We now define an Ornstein-Uhlenbeck process on $C(M)$ of covariance C_μ and drift $-G_\mu$. This heavily relies on the general construction given in the appendix.

A *Brownian motion on $C(M)$ with covariance C_μ* is a $C(M)$ -valued stochastic process $W = \{W_t\}_{t \geq 0}$ such that

- (i) $W_0 = 0$;
- (ii) $t \mapsto W_t$ is continuous;
- (iii) For every finite subset $S \subset \mathbb{R} \times M$, $\{W_t(x)\}_{(t,x) \in S}$ is a centered Gaussian random vector;
- (iv) $\mathbb{E}[W_s(x)W_t(y)] = (s \wedge t)C_\mu(x, y)$.

Lemma 2.3 *There exists a Brownian motion on $C(M)$ with covariance C_μ .*

Proof : Let

$$\begin{aligned} d_{C_\mu}(x, y) &:= \sqrt{C_\mu(x, x) - 2C_\mu(x, y) + C_\mu(y, y)} \\ &= \|\nabla \mathbf{Q}_\mu(V_x - V_y)\|_{\Pi(\mu)} \\ &\leq K\|V_x - V_y\| \end{aligned}$$

where the last inequality follows from (9). Then

$$d_{C_\mu}(x, y) \leq Kd(x, y)$$

and the result follows from Proposition 5.8 and Remark 5.7 in the appendix.

QED

We say that a $C(M)$ -valued process Z is an *Ornstein-Uhlenbeck process of covariance C_μ and drift $-G_\mu$* if

$$Z_t = Z_0 - \int_0^t G_\mu Z_s ds + W_t \quad (11)$$

where

- (i) W is a $C(M)$ -valued Brownian motion of covariance C_μ ;
- (ii) Z_0 is a $C(M)$ -valued random variable;
- (iii) W and Z_0 are independent.

Note that we can think of Z as a solution to the linear SDE

$$dZ_t = dW_t - G_\mu Z_t dt.$$

It follows from section 5.3 in the appendix that such a process exists and defines a Markov process. Furthermore

Proposition 2.4 *Under hypothesis 2.1,*

- (i) (Z_t) converges in law toward a $C(M)$ -valued random variable Z_∞ ;
- (ii) Z_∞ is Gaussian, in the sense that for every finite set $S \subset M$, $\{Z_\infty(x)\}_{x \in S}$ is a centered Gaussian random vector;
- (iii) Let π^μ denotes the law of Z_∞ . Then π^g is characterized by its variance

$$\text{Var}(\pi^\mu) : \mathcal{M}(M) \rightarrow \mathbb{R},$$

$$m \mapsto \mathbf{E}((mZ_\infty)^2),$$

and for all $m \in \mathcal{M}$,

$$\begin{aligned}\text{Var}(\pi^\mu)(m) &= \int_0^\infty \int_{M \times M} C_\mu(x, y) m_t(dx) m_t(dy) dt \\ &= \int_0^\infty \widehat{C}_\mu(V m_t, V m_t) dt\end{aligned}$$

where

$$m_t = e^{-tG_\mu^*} m.$$

Proof : This follows from Proposition 5.16 in the appendix. Example 5.18 shows that assertion (iii) of this proposition is satisfied. **QED**

2.5 The process Z^g .

For $g = (g_1, \dots, g_n) \in C(M)^n$, let $\tilde{M} = \{1, \dots, n\} \cup M$ be the disjoint union of $\{1, \dots, n\}$ and M , and $C_\mu^g : \tilde{M} \times \tilde{M} \rightarrow \mathbb{R}$ be the function defined by

$$C_\mu^g(x, y) = \begin{cases} \widehat{C}_\mu(g_x, g_y) & \text{for } x, y \in \{1, \dots, n\}, \\ C_\mu(x, y) & \text{for } x, y \in M, \\ \widehat{C}_\mu(V_x, g_y) & \text{for } x \in M, y \in \{1, \dots, n\}. \end{cases}$$

Then C_μ^g is a Mercer kernel (see section 5.2).

A *Brownian motion on $\mathbb{R}^n \times C(M)$ with covariance C_μ^g* is a $\mathbb{R}^n \times C(M)$ -valued stochastic process $(W^g, W) = \{(W_t^{g_1}, \dots, W_t^{g_n}, W_t)\}_{t \geq 0}$ such that:

- (i) $W = \{W_t\}_{t \geq 0}$ is a $C(M)$ -valued Brownian motion with covariance C_μ ;
- (ii) For every finite subset $S \subset \mathbb{R} \times M$, $\{W_t^g, W_t(x)\}_{(t,x) \in S}$ is a centered Gaussian random vector;
- (iii) $E(W_s^{g_i} W_t^{g_j}) = (s \wedge t) \widehat{C}_\mu(g_i, g_j)$ and $E(W_s(x) W_t^{g_i}) = (s \wedge t) \widehat{C}_\mu(V_x, g_i)$.

Lemma 2.5 *There exists a Brownian motion on $\mathbb{R}^n \times C(M)$ with covariance C_μ^g .*

Proof : Let $\tilde{d}$ be the distance on $\tilde{M}$ defined by

$$\tilde{d}(x, y) = \begin{cases} \mathbf{1}_{x \neq y} & \text{for } x, y \in \{1, \dots, n\}, \\ d(x, y) & \text{for } x, y \in M, \\ d(x, x_0) + 1 & \text{for } x \in M, y \in \{1, \dots, n\} \end{cases}$$

where x_0 is some arbitrary point in M . This makes $\tilde{M}$ a compact metric space, and it is easy to show that the function C_μ^g verifies hypothesis 5.6 (use the proof of Lemma 2.3). The result follows by application of Proposition 5.8. **QED**

Let now be $Z_t^g = (Z_t^{g_1}, \dots, Z_t^{g_n}) \in \mathbb{R}^n$ denote the solution to the SDE

$$dZ_t^{g_i} = dW_t^{g_i} - (Z_t^{g_i}/2 + \text{Cov}_\mu(Z_t, g_i)) dt, \quad i = 1, \dots, n \quad (12)$$

where (W^g, W) is as above and $Z = (Z_t)$ is given by (11).

The following result generalizes Proposition 2.4.

Proposition 2.6 *Under hypothesis 2.1,*

- (i) *The process (Z_t^g, Z_t) converges in law toward a centered $\mathbb{R}^n \times C(M)$ valued Gaussian random variable (Z_∞^g, Z_∞) .*
- (ii) *Let $\pi^{g,\mu}$ denotes the law of (Z_∞^g, Z_∞) . Then $\pi^{g,\mu}$ is characterized by its variance*

$$\text{Var}(\pi^{g,\mu}) : \mathbb{R}^n \times \mathcal{M}(M) \rightarrow \mathbb{R},$$

$$(u, m) \mapsto \mathbb{E}((mZ_\infty + \langle u, Z_\infty^g \rangle)^2);$$

and for all $u \in \mathbb{R}^n, m \in \mathcal{M}(M)$,

$$\text{Var}(\pi^{g,\mu})(u, m) = \int_0^\infty \widehat{C}_\mu(f_t, f_t) dt$$

with

$$f_t = e^{-t/2} \sum_i u_i g_i + V m_t,$$

and where m_t is defined by

$$m_t f = m_0(e^{-tG_\mu} f) + \sum_{i=1}^n u_i \int_0^t e^{-s/2} \text{Cov}_\mu(g_i, e^{-(t-s)G_\mu} f) ds. \quad (13)$$

Proof : Let $G_\mu^g : \mathbb{R}^n \times C(M) \rightarrow \mathbb{R}^n \times C(M)$ be the operator defined by

$$G_\mu^g = \begin{pmatrix} I/2 & A_\mu^g \\ 0 & G_\mu \end{pmatrix} \quad (14)$$

where $A_\mu^g : C(M) \rightarrow \mathbb{R}^n$ is the linear map defined by

$$A_\mu^g(f) = \left(\text{Cov}_\mu(f, g_1), \dots, \text{Cov}_\mu(f, g_n) \right).$$

Then (Z^g, Z) is a $C(\tilde{M})$ -valued Ornstein-Uhlenbeck process of covariance C_μ^g and drift $-G_\mu^g$. It is not hard to verify that under hypothesis 2.1, the assumptions of Proposition 5.16 hold, so that (Z_t^g, Z_t) converges in law toward a centered $\mathbb{R}^n \times C(M)$ valued Gaussian random variable (Z_∞^g, Z_∞) with variance

$$\text{Var}(\pi^{g,\mu})(u, m) = \int_0^\infty \widehat{C}_\mu(f_t, f_t) dt$$

with $f_t = \sum_i u_t(i)g_i + Vm_t$ and where $(u_t, m_t) = e^{-t(G_\mu^g)^*}(u, m)$. Now

$$(G_\mu^g)^* = \begin{pmatrix} I/2 & 0 \\ (A_\mu^g)^* & (G_\mu)^* \end{pmatrix}$$

and $(A_\mu^g)^*u = \sum_i u_i(g_i - \mu g_i)\mu$. Thus $u_t = e^{-t/2}u$ and

$$\frac{dm_t}{dt} = -(A_\mu^g)^*u_t - (G_\mu)^*m_t$$

Thus m_t is the solution with $m_0 = m$ of

$$\frac{dm_t}{dt} = -e^{-t/2} \left(\sum_i u_i(g_i - \mu g_i) \right) \mu - G_\mu^* m_t \quad (15)$$

Note that (15) is equivalent to

$$\frac{d}{dt}(m_t f) = -e^{-t/2} \text{Cov}_\mu \left(\sum_i u_i g_i, f \right) - m_t (G_\mu f)$$

for all $f \in C(M)$, and $m_0 = m$. From which we deduce that

$$m_t = e^{-tG_\mu^*} m_0 - \int_0^t e^{-s/2} e^{-(t-s)G_\mu^*} \left(\sum_i u_i(g_i - \mu g_i) \mu \right) ds$$

which implies the formula for m_t given by (13). **QED**

For further reference we call (Z^g, Z) an *Ornstein-Uhlenbeck process of covariance C_μ^g and drift $-G_\mu^g$* . It is called *stationary* when its initial distribution is $\pi^{g,\mu}$.

3 A central limit theorem for μ_t

We state here the main results of this article. We assume $\mu^* \in \text{Fix}(\Pi)$ satisfies hypotheses 1.3 and 2.1. Set $\Delta_t = e^{t/2}(\mu_{e^t} - \mu^*)$, $D_t = V\Delta_t$ and $D_{t+} = \{D_{t+s} : s \geq 0\}$. Then

Theorem 3.1 *D_{t+} converges in law, as $t \rightarrow \infty$, towards a stationary Ornstein-Uhlenbeck process of covariance C_{μ^*} and drift $-G_{\mu^*}$.*

For $g = (g_1, \dots, g_n) \in C(M)^n$, we set $D_t^g = (\Delta_t g, D_t)$ and $D_{t+}^g = \{D_{t+s}^g : s \geq 0\}$. Then

Theorem 3.2 *$(D_{t+s}^g)_{s \geq 0}$ converges in law towards a stationary Ornstein-Uhlenbeck process of covariance $C_{\mu^*}^g$ and drift $-G_{\mu^*}^g$.*

Define $\widehat{C} : C(M) \times C(M) \rightarrow \mathbb{R}$ the symmetric bilinear form defined by

$$\widehat{C}(f, g) = \int_0^\infty \widehat{C}_{\mu^*}(f_t, g_t) dt, \quad (16)$$

with (g_t) is defined by the same formula, with g in place of f)

$$f_t(x) = e^{-t/2} f(x) - \int_0^t e^{-s/2} \text{Cov}_{\mu^*}(f, e^{-(t-s)G_{\mu^*}} V_x) ds. \quad (17)$$

Corollary 3.3 *$\Delta_t g$ converges in law towards a centered Gaussian variable Z_∞^g of covariance*

$$\mathbb{E}[Z_\infty^{g_i} Z_\infty^{g_j}] = \widehat{C}(f, g).$$

Proof : Follows from theorem 3.2 and the calculus of $\text{Var}(\pi^{g, \mu})(u, 0)$. **QED**

3.1 Examples

3.1.1 Diffusions

Suppose $V(x, y) = V(x)$, so that (X_t) is just a standard diffusion on M with invariant measure $\mu^* = \frac{\exp(-V)\lambda}{\lambda \exp(-V)}$.

Let $f \in C(M)$. Then f_t defined by (17) is equal to (using $e^{-tG_{\mu^*}} 1 = e^{-t/2} 1$) $= e^{-t/2} f$. Thus

$$\widehat{C}(f, g) = 2\mu^*(f Q_{\mu^*} g). \quad (18)$$

Corollary 3.3 says that

Theorem 3.4 *For all $g \in C(M)^n$, Δ_t^g converges in law toward a centered Gaussian variable $(Z_\infty^{g_1}, \dots, Z_\infty^{g_n})$, with covariance given by*

$$\mathbb{E}(Z_\infty^{g_i} Z_\infty^{g_j}) = 2\mu^*(g_i \mathbf{Q}_{\mu^*} g_j).$$

Remark 3.5 *This central limit theorem for Brownian motions on compact manifolds has already been considered by Baxter and Brosamer in [1] and [2]; and by Bhattacharya in [7] for ergodic diffusions.*

3.1.2 The case $\mu^* = \lambda$ and V symmetric.

Suppose here that $\mu^* = \lambda$ and that V is symmetric. We assume (without loss of generality since $\Pi(\lambda) = \lambda$ implies that $V\lambda$ is a constant function) that $V\lambda = 0$.

Since V is compact and symmetric, there exists an orthonormal basis $(e_\alpha)_{\alpha \geq 0}$ in $L^2(\lambda)$ and a sequence of reals $(\lambda_\alpha)_{\alpha \geq 0}$ such that e_0 is a constant function and

$$V = \sum_{\alpha \geq 1} \lambda_\alpha e_\alpha \otimes e_\alpha.$$

Assume that for all α , $1/2 + \lambda_\alpha > 0$. Then hypothesis 2.1 holds with $\hat{\lambda} = \lambda$, and the convergence of μ_t towards λ holds with positive probability (see [6]).

Let $f \in C(M)$ and f_t defined by (17), denoting $f^\alpha = \langle f, e_\alpha \rangle_\lambda$ and $f_t^\alpha = \langle f_t, e_\alpha \rangle_\lambda$, we have $f_t^0 = e^{-t/2} f^0$ and for $\alpha \geq 1$,

$$\begin{aligned} f_t^\alpha &= e^{-t/2} f^\alpha - \lambda_\alpha e^{-(1/2 + \lambda_\alpha)t} \left(\frac{e^{\lambda_\alpha t} - 1}{\lambda_\alpha} \right) f^\alpha \\ &= e^{-(1/2 + \lambda_\alpha)t} f^\alpha. \end{aligned}$$

Using the fact that

$$\hat{C}_\lambda(f, g) = 2\lambda(f \mathbf{Q}_\lambda g),$$

this implies that

$$\hat{C}(f, g) = 2 \sum_{\alpha \geq 1} \sum_{\beta \geq 1} \frac{1}{1 + \lambda_\alpha + \lambda_\beta} \langle f, e_\alpha \rangle_\lambda \langle g, e_\beta \rangle_\lambda \lambda(e_\alpha \mathbf{Q}_\lambda e_\beta).$$

This, with corollary 3.3, proves

Theorem 3.6 *Assume hypothesis 1.3 and that $1/2 + \lambda_\alpha > 0$ for all α . Then for all $g \in C(M)^n$, Δ_t^g converges in law toward a centered Gaussian variable $(Z_\infty^{g_1}, \dots, Z_\infty^{g_n})$, with covariance given by*

$$\mathbb{E}(Z_\infty^{g_i} Z_\infty^{g_j}) = \widehat{C}(g_i, g_j).$$

In particular,

$$\mathbb{E}(Z_\infty^{e_\alpha} Z_\infty^{e_\beta}) = \frac{2}{1 + \lambda_\alpha + \lambda_\beta} \lambda(e_\alpha \mathbf{Q}_\lambda e_\beta).$$

Note that when all λ_α are positive, which corresponds to what is named a self-repelling interaction in [6], the rate of convergence of μ_t towards λ is bigger than when there is no interaction, and the bigger is the interaction (that is larger λ_α 's) faster is the convergence.

4 Proof of the main results

We assume hypothesis 1.3 and μ^* satisfies hypothesis 2.1. It is possible to choose κ in hypothesis 2.1 such that $\kappa < 1/2$. In the following κ will denote such constant. Note that we have $\lambda(-G_{\mu^*}) < -\kappa$. Such κ exists when hypothesis 2.1 holds.

4.1 A lemma satisfied by $\mathbf{Q}_\mu$

We denote by $\mathcal{X}(M)$ the space of continuous vector fields on M , and equip the spaces $\mathcal{P}(M)$ and $\mathcal{X}(M)$ respectively with the weak convergence topology and with the uniform convergence topology.

Lemma 4.1 *For all $f \in C(M)$, the mapping $\mu \mapsto \nabla \mathbf{Q}_\mu f$ is a continuous mapping from $\mathcal{P}(M)$ in $\mathcal{X}(M)$.*

Proof : Let μ and ν be in $\mathcal{M}(M)$, and $f \in C(M)$. Set $g = \mathbf{Q}_\mu f$. Then $f = -A_\mu g + \Pi(\mu)f$ and

$$\begin{aligned} \|\nabla \mathbf{Q}_\mu f - \nabla \mathbf{Q}_\nu f\|_\infty &= \|\nabla \mathbf{Q}_\mu f - \nabla \mathbf{Q}_\mu A_\mu g + \nabla \mathbf{Q}_\nu A_\mu g\|_\infty \\ &= \|\nabla g + \nabla \mathbf{Q}_\nu A_\mu g\|_\infty \\ &\leq \|\nabla(g + \mathbf{Q}_\nu A_\nu g)\|_\infty + \|\nabla \mathbf{Q}_\nu(A_\mu - A_\nu)g\|_\infty \end{aligned}$$

since $\nabla(g + Q_\nu A_\nu g) = 0$ and $(A_\mu - A_\nu)g = \langle \nabla V_{\mu-\nu}, \nabla g \rangle$, we get

$$\|\nabla Q_\mu f - \nabla Q_\nu f\|_\infty \leq K \|\langle \nabla V_{\mu-\nu}, \nabla g \rangle\|_\infty. \quad (19)$$

Using the fact that $(x, y) \mapsto \nabla V_x(y)$ is uniformly continuous, the right hand term of (19) converges towards 0, when $d(\mu, \nu)$ converges towards 0, d being a distance compatible with the weak convergence. **QED**

4.2 The process Δ

Set $h_t = V\mu_t$ and $h^* = V\mu^*$. Recall $\Delta_t = e^{t/2}(\mu_{e^t} - \mu^*)$ and $D_t = V\Delta_t$. Note that $D_t(x) = \Delta_t V_x$.

To simplify the notation, we set $K_s = K_{\mu_s}$, $Q_s = Q_{\mu_s}$ and $A_s = A_{\mu_s}$. Let $(M_t^f)_{t \geq 1}$ be the martingale defined by

$$M_t^f = \sum_{\alpha} \int_1^t e_{\alpha}(Q_s f)(X_s) dB_s^{\alpha}.$$

The quadratic covariation of M^f and M^g (with f and g in $C(M)$) is given by

$$\langle M^f, M^g \rangle_t = \int_1^t \langle \nabla Q_s f, \nabla Q_s g \rangle(X_s) ds.$$

Then for all $t \geq 1$ (with $\dot{Q}_t = \frac{d}{dt} Q_t$),

$$Q_t f(X_t) - Q_1 f(X_1) = M_t^f + \int_1^t \dot{Q}_s f(X_s) ds - \int_1^t K_s f(X_s) ds.$$

Thus

$$\begin{aligned} \mu_t f &= \frac{1}{t} \int_1^t K_s f(X_s) ds + \frac{1}{t} \int_1^t \Pi(\mu_s) f ds + \frac{1}{t} \int_0^1 f(X_s) ds \\ &= -\frac{1}{t} \left(Q_t f(X_t) - Q_1 f(X_1) - \int_1^t \dot{Q}_s f(X_s) ds \right) \\ &\quad + \frac{M_t^f}{t} + \frac{1}{t} \int_1^t \langle \xi(h_s), f \rangle_{\lambda} ds + \frac{1}{t} \int_0^1 f(X_s) ds. \end{aligned}$$

Note that (D_t) is a continuous process taking its values in $C(M)$ and that $D_t = e^{t/2}(h_{e^t} - h^*)$. For $f \in C(M)$ (using the fact that $\mu^* f = \langle \xi(h^*), f \rangle_{\lambda}$),

$$\Delta_t f = \sum_{i=1}^5 \Delta_t^i f \quad (20)$$

with

$$\begin{aligned}
\Delta_t^1 f &= e^{-t/2} \left(-\mathbf{Q}_{e^t} f(X_{e^t}) + \mathbf{Q}_1 f(X_1) + \int_1^{e^t} \dot{\mathbf{Q}}_s f(X_s) ds \right) \\
\Delta_t^2 f &= e^{-t/2} M_{e^t}^f \\
\Delta_t^3 f &= e^{-t/2} \int_1^{e^t} \langle \xi(h_s) - \xi(h^*) - D\xi(h^*)(h_s - h^*), f \rangle_\lambda ds \\
\Delta_t^4 f &= e^{-t/2} \int_1^{e^t} \langle D\xi(h^*)(h_s - h^*), f \rangle_\lambda ds \\
\Delta_t^5 f &= e^{-t/2} \left(\int_0^1 f(X_s) ds - \mu^* f \right).
\end{aligned}$$

Then $D_t = \sum_{i=1}^5 D_t^i$, where $D_t^i = V \Delta_t^i$. Finally, note that

$$\langle D\xi(h^*)(h - h^*), f \rangle_\lambda = -\text{Cov}_{\mu^*}(h - h^*, f). \quad (21)$$

4.3 First estimates

We recall some estimates from [3]: There exists a constant K such that for all $f \in C(M)$ and $t > 0$,

$$\begin{aligned}
\|\mathbf{Q}_t f\|_\infty &\leq K \|f\|_\infty \\
\|\nabla \mathbf{Q}_t f\|_\infty &\leq K \|f\|_\infty \\
\|\dot{\mathbf{Q}}_t f\|_\infty &\leq \frac{K}{t} \|f\|_\infty.
\end{aligned}$$

These estimates imply in particular that

$$\langle M^f - M^g \rangle_t \leq K \|f - g\|_\infty \times t$$

and that

Lemma 4.2 *There exists a constant K depending on $\|V\|_\infty$ such that for all $t \geq 1$, and all $f \in C(M)$*

$$\|\Delta_t^1 f\|_\infty + \|\Delta_t^5 f\|_\infty \leq K \times (1 + t) e^{-t/2} \|f\|_\infty, \quad (22)$$

which implies that $((\Delta^1 + \Delta^5)_{t+s})_{s \geq 0}$ and $((D^1 + D^5)_{t+s})_{s \geq 0}$ both converge towards 0 (respectively in $\mathcal{M}(M)$ and in $C(\mathbb{R}^+ \times M)$).

We also have

Lemma 4.3 *There exists a constant K such that for all $t \geq 0$ and all $f \in C(M)$,*

$$\begin{aligned} \mathbb{E}[(\Delta_t^2 f)^2] &\leq K \|f\|_\infty^2, \\ |\Delta_t^3 f| &\leq K \|f\|_\lambda \times e^{-t/2} \int_0^t \|D_s\|_\lambda^2 ds, \\ |\Delta_t^4 f| &\leq K \|f\|_\lambda \times e^{-t/2} \int_0^t e^{s/2} \|D_s\|_\lambda ds. \end{aligned}$$

Proof : The first estimate follows from

$$\begin{aligned} \mathbb{E}[(\Delta_t^2 f)^2] &= e^{-t} \mathbb{E}[(M_{e^t}^f)^2] = e^{-t} \mathbb{E}[\langle M^f \rangle_{e^t}] \\ &\leq e^{-t} \int_1^{e^t} \|\nabla Q_s f\|_\infty^2 ds \\ &\leq K \|f\|_\infty^2. \end{aligned}$$

The second estimate follows from the fact that

$$\|\xi(h) - \xi(h^*) - D\xi(h^*)(h - h^*)\|_\lambda = O(\|h - h^*\|_\lambda^2).$$

The last estimate follows easily after having remarked that

$$\begin{aligned} |\langle D\xi(h^*)(h_s - h^*), f \rangle| &= |\text{Cov}_{\mu^*}(h_s - h^*, f)| \\ &\leq K \|f\|_\lambda \times \|h_s - h^*\|_\lambda \\ &\leq K \|f\|_\lambda \times s^{-1/2} \|D_{\log(s)}\|_\lambda. \end{aligned}$$

This proves this lemma. **QED**

4.4 The processes Δ' and D'

Set $\Delta' = \Delta^2 + \Delta^3 + \Delta^4$ and $D' = D^2 + D^3 + D^4$. For $g \in C(M)$, set

$$\epsilon_t^g = e^{t/2} \langle \xi(h_{e^t}) - \xi(h^*) - D\xi(h^*)(h_{e^t} - h^*), g \rangle_\lambda.$$

Then

$$d\Delta'_t g = -\frac{\Delta'_t g}{2} dt + dN_t^g + \epsilon_t^g dt + \langle D\xi(h^*)(D_t), g \rangle_\lambda dt$$

where for all $g \in C(M)$, N^g is a martingale. Moreover, for f and g in $C(M)$,

$$\langle N^f, N^g \rangle_t = \int_0^t \langle \nabla Q_{e^s} f(X_{e^s}), \nabla Q_{e^s} g(X_{e^s}) \rangle ds.$$

Then, for all x ,

$$dD'_t(x) = -\frac{D'_t(x)}{2}dt + dM_t(x) + \epsilon_t(x)dt + \langle D\xi(h^*)(D_t), V_x \rangle_\lambda dt$$

where M is the martingale in $C(M)$ defined by $M(x) = N^{V_x}$ and $\epsilon_t(x) = \epsilon_t^{V_x}$. We also have

$$G_{\mu^*}(D')_t(x) = \frac{D'_t(x)}{2} - \langle D\xi(h^*)(D'_t), V_x \rangle_\lambda.$$

Denoting $L_{\mu^*} = L_{-G_{\mu^*}}$ (defined by equation (32) in the appendix), this implies that

$$\begin{aligned} dL_{\mu^*}(D')_t(x) &= dD'_t(x) + G_{\mu^*}(D')_t(x)dt \\ &= dM_t(x) + \langle D\xi(h^*)((D^1 + D^5)_t), V_x \rangle_\lambda dt + \epsilon_t(x)dt \end{aligned}$$

Thus

$$L_{\mu^*}(D')_t(x) = M_t(x) + \int_0^t \epsilon'_s(x)ds$$

with $\epsilon'_s(x) = \epsilon'_s V_x$ where for all $f \in C(M)$,

$$\epsilon'_s f = \epsilon_s^f + \langle D\xi(h^*)((D^1 + D^5)_s), f \rangle_\lambda.$$

Using lemma 5.10,

$$D'_t = L_{\mu^*}^{-1}(M)_t + \int_0^t e^{-(t-s)G_{\mu^*}} \epsilon'_s ds. \quad (23)$$

For $g = (g_1, \dots, g_n) \in C(M)^n$, we denote $\Delta'_t g = (\Delta'_t g_1, \dots, \Delta'_t g_n)$, $N^g = (N^{g_1}, \dots, N^{g_n})$ and $\epsilon'_t g = (\epsilon'_t g_1, \dots, \epsilon'_t g_n)$. Then, denoting $L_{\mu^*}^g = L_{-G_{\mu^*}^g}$ (with $G_{\mu^*}^g$ defined by (14)) we have

$$L_{\mu^*}^g(\Delta'_t g, D')_t = (N_t^g, M_t) + \int_0^t (\epsilon'_s g, \epsilon'_s) ds$$

so that (using lemma 5.10 and integrating by parts)

$$(\Delta'_t g, D'_t) = (L_{\mu^*}^g)^{-1}(N^g, M)_t + \int_0^t e^{-(t-s)G_{\mu^*}^g}(\epsilon'_s g, \epsilon'_s) ds. \quad (24)$$

Moreover

$$(L_{\mu^*}^g)^{-1}(N^g, M)_t = \left(\widehat{N}_t^{g_1}, \dots, \widehat{N}_t^{g_n}, L_{\mu^*}^{-1}(M)_t \right),$$

where

$$\widehat{N}_t^{g_i} = N_t^{g_i} - \int_0^t \left(\frac{N_s^{g_i}}{2} + \widehat{C}_{\mu^*}(L_{\mu^*}^{-1}(M)_s, g_i) \right) ds.$$

4.5 Estimation of ϵ'_t

4.5.1 Estimation of $\|L_{\mu^*}^{-1}(M)_t\|_\lambda$

Lemma 4.4 (i) *For all $\alpha \geq 2$, there exists a constant K_α such that for all $t \geq 0$,*

$$\mathbb{E}[\|L_{\mu^*}^{-1}(M)_t\|_\lambda^\alpha]^{1/\alpha} \leq K_\alpha.$$

(ii) *a.s. there exists C with $\mathbb{E}[C] < \infty$ such that for all $t \geq 0$,*

$$\|L_{\mu^*}^{-1}(M)_t\|_\lambda \leq C(1+t).$$

Proof : Since $\|L_{\mu^*}^{-1}(M)_t\|_\lambda \leq K\|L_{\mu^*}^{-1}(M)_t\|_{\widehat{\lambda}}$, we estimate $\|L_{\mu^*}^{-1}(M)_t\|_{\widehat{\lambda}}$. We have

$$dL_{\mu^*}^{-1}(M)_t = dM_t - G_{\mu^*} L_{\mu^*}^{-1}(M)_t dt.$$

Let N be the martingale defined by

$$N_t = \int_0^t \left\langle \frac{L_{\mu^*}^{-1}(M)_s}{\|L_{\mu^*}^{-1}(M)_s\|_{\widehat{\lambda}}}, dM_s \right\rangle_{\widehat{\lambda}}.$$

We have $\langle N \rangle_t \leq Kt$ for some constant K . Then

$$\begin{aligned} d\|L_{\mu^*}^{-1}(M)_t\|_{\widehat{\lambda}}^2 &= 2\|L_{\mu^*}^{-1}(M)_t\|_{\widehat{\lambda}} dN_t - 2\langle L_{\mu^*}^{-1}(M)_t, G_{\mu^*} L_{\mu^*}^{-1}(M)_t \rangle_{\widehat{\lambda}} dt \\ &\quad + d\left(\int \langle M(x) \rangle_t \widehat{\lambda}(dx)\right). \end{aligned}$$

Note that there exists a constant K such that

$$\frac{d}{dt} \left(\int \langle M(x) \rangle_t \widehat{\lambda}(dx) \right) \leq K$$

and that (see hypothesis 2.1)

$$\langle L_{\mu^*}^{-1}(M)_t, G_{\mu^*} L_{\mu^*}^{-1}(M)_t \rangle_{\hat{\lambda}} \geq \kappa \|L_{\mu^*}^{-1}(M)_t\|_{\hat{\lambda}}^2.$$

This implies that

$$\frac{d}{dt} \mathbb{E}[\|L_{\mu^*}^{-1}(M)_t\|_{\hat{\lambda}}^2] \leq -2\kappa \mathbb{E}[\|L_{\mu^*}^{-1}(M)_t\|_{\hat{\lambda}}^2] + K$$

which implies (i) for $\alpha = 2$. For $\alpha > 2$, we find that

$$\begin{aligned} \frac{d}{dt} \mathbb{E}[\|L_{\mu^*}^{-1}(M)_t\|_{\hat{\lambda}}^\alpha] &\leq -\alpha\kappa \mathbb{E}[\|L_{\mu^*}^{-1}(M)_t\|_{\hat{\lambda}}^\alpha] + K \mathbb{E}[\|L_{\mu^*}^{-1}(M)_t\|_{\hat{\lambda}}^{\alpha-2}] \\ &\leq -\alpha\kappa \mathbb{E}[\|L_{\mu^*}^{-1}(M)_t\|_{\hat{\lambda}}^\alpha] + K \mathbb{E}[\|L_{\mu^*}^{-1}(M)_t\|_{\hat{\lambda}}^\alpha]^{\frac{\alpha-2}{\alpha}} \end{aligned}$$

which implies that $\mathbb{E}[\|L_{\mu^*}^{-1}(M)_t\|_{\hat{\lambda}}^\alpha]$ is bounded.

We now prove (ii). Fix $\alpha > 1$. Then there exists a constant K such that

$$\frac{\|L_{\mu^*}^{-1}(M)_t\|_{\hat{\lambda}}^2}{(1+t)^\alpha} \leq \|L_{\mu^*}^{-1}(M)_0\|_{\hat{\lambda}}^2 + 2 \int_0^t \frac{\|L_{\mu^*}^{-1}(M)_s\|_{\hat{\lambda}}}{(1+s)^\alpha} dN_s + K.$$

Then BDG inequality implies that

$$\mathbb{E} \left[\sup_{t \geq 0} \frac{\|L_{\mu^*}^{-1}(M)_t\|_{\hat{\lambda}}^2}{(1+t)^\alpha} \right] \leq K + 2 \sup_{t \geq 0} \left(\int_0^t \frac{K ds}{(1+s)^{2\alpha}} \right)^{1/2}$$

which is finite. This implies the lemma by taking $\alpha = 2$. **QED**

4.5.2 Estimation of $\|D_t\|_\lambda$

Note that $|\epsilon_t^g| \leq K e^{-t/2} \|D_t\|_\lambda^2 \times \|g\|$. Thus

$$|\epsilon_t' g| \leq K e^{-t/2} (1+t+\|D_t\|_\lambda^2) \times \|g\|.$$

This implies (using lemma 2.2 and the fact that $0 < \kappa < 1/2$)

Lemma 4.5 *There exists K such that*

$$\left\| \int_0^t e^{-(t-s)G_{\mu^*}} \epsilon_s' ds \right\| \leq K e^{-\kappa t} \left(1 + \int_0^t e^{-(1/2-\kappa)s} \|D_s\|_\lambda^2 ds \right). \quad (25)$$

This lemma with lemma 4.4-(ii) imply the following

Lemma 4.6 *a.s. there exists C with $\mathbf{E}[C] < \infty$ such that*

$$\|D_t\|_\lambda \leq C \times \left[1 + t + \int_0^t e^{-s/2} \|D_s\|_\lambda^2 ds \right]. \quad (26)$$

Proof : First note that

$$\|D_t\|_\lambda \leq \|D'_t\|_\lambda + K(1+t)e^{-t/2}.$$

Using the expression of D'_t given by (23), we get

$$\begin{aligned} \|D'_t\|_\lambda &\leq \|L_{\mu^*}^{-1}(M)_t\|_\lambda + \left\| \int_0^t e^{-(t-s)G_{\mu^*}} \epsilon'_s ds \right\| \\ &\leq C(1+t) + Ke^{-\kappa t} \left(1 + \int_0^t e^{-(1/2-\kappa)s} \|D_s\|_\lambda^2 ds \right) \end{aligned}$$

which implies the lemma. **QED**

Lemma 4.7 *Let x and ϵ be real functions. If for all $t \geq 0$,*

$$x_t \leq \alpha + \int_0^t \epsilon_s x_s ds,$$

where α is a real constant, then

$$x_t \leq \alpha \exp \left(\int_0^t \epsilon_s ds \right).$$

Proof : Similarly to the proof of Gronwall's lemma, we set $y_t = \int_0^t \epsilon_s x_s ds$. Then,

$$\dot{y}_t \leq \alpha \epsilon_t + \epsilon_t y_t.$$

Take $\lambda_t = y_t \exp \left(- \int_0^t \epsilon_s ds \right)$, then

$$\dot{\lambda}_t \leq \alpha \epsilon_t \exp \left(- \int_0^t \epsilon_s ds \right)$$

and

$$\begin{aligned} y_t &\leq \alpha \int_0^t \epsilon_s \exp \left(\int_s^t \epsilon_u du \right) ds \\ &\leq \alpha \exp \left(\int_0^t \epsilon_u du \right) - \alpha. \end{aligned}$$

This implies the lemma. **QED**

This lemma implies that

$$\|D_t\|_\lambda \leq C(1+t) \times \exp \left(C \int_0^t e^{-s/2} \|D_s\|_\lambda ds \right).$$

Since hypothesis 1.3 implies that $\lim_{s \rightarrow \infty} e^{-s/2} \|D_s\|_\lambda = 0$, this proves that a.s. for all $\epsilon > 0$, there exists C_ϵ such that

$$\|D_t\|_\lambda \leq C_\epsilon e^{\epsilon t}.$$

Take $\epsilon < 1/4$. Then

$$\int_0^\infty e^{-s/2} \|D_s\|_\lambda^2 ds \leq C_\epsilon.$$

This implies

Lemma 4.8 *a.s., there exists C such that for all t ,*

$$\|D_t\|_\lambda \leq C(1+t).$$

4.5.3 Estimation of ϵ'_t

Lemma 4.9 *a.s. there exists C such that for all $f \in C(M)$,*

$$|\epsilon'_t f| \leq C(1+t)^2 e^{-t/2} \|f\|$$

Proof : We have $|\epsilon'_t f| \leq |\epsilon_t^f| + K(1+t)e^{-t/2} \|f\|$ and

$$\begin{aligned} |\epsilon_t^f| &\leq K \|f\|_\lambda \times e^{-t/2} \|D_t\|_\lambda^2 \\ &\leq C \|f\| \times (1+t)^2 e^{-t/2} \end{aligned}$$

by lemma 4.8. **QED**

4.6 Estimation of $\|D_t - L_{\mu^*}^{-1}(M)_t\|$

Lemma 4.10 $\|D_t - L_{\mu^*}^{-1}(M)_t\| \leq Ce^{-\kappa t}$.

Proof : We have $\|D_t - D'_t\| \leq K(1+t)e^{-t/2}$. So to prove this lemma, it suffices to prove that (see the expression of D'_t given by (23))

$$\left\| \int_0^t e^{-(t-s)G_{\mu^*}} \epsilon'_s ds \right\| \leq Ce^{-\kappa t}.$$

This term is dominated by

$$K \int_0^t e^{-\kappa(t-s)} \|\epsilon'_s\| ds.$$

Using the previous lemma, it is also dominated by

$$Ce^{-\kappa t} \int_0^t e^{\kappa s} (1+s)^2 e^{-s/2} ds \leq Ce^{-\kappa t}$$

because $\kappa \in]0, 1/2[$. The lemma is proved. **QED**

In addition, for $g = (g_1, \dots, g_n) \in C(M)^n$, setting

$$\Delta_t g = (\Delta_t g_1, \dots, \Delta_t g_n),$$

Lemma 4.11 $\|(\Delta_t g, D_t) - (L_{\mu^*}^g)^{-1}(N^g, M)_t\| \leq C(1 + \|g\|)e^{-\kappa t}$.

Proof : We have $\|(\Delta_t g, D_t) - (\Delta'_t g, D'_t)\| \leq K(1 + \|g\|)(1+t)e^{-\kappa t}$. So to prove this lemma, using (24), it suffices to prove that

$$\left\| \int_0^t e^{-(t-s)G_{\mu^*}^g} (\epsilon'_s g, \epsilon'_s) ds \right\| \leq K(1 + \|g\|)e^{-\kappa t}. \quad (27)$$

Using hypothesis 2.1 and the definition of $G_{\mu^*}^g$, we have that for all positive t ,

$$\|e^{-tG_{\mu^*}^g}\| \leq Ke^{-\kappa t}.$$

This implies

$$\|e^{-(t-s)G_{\mu^*}^g} (\epsilon'_s g, \epsilon'_s)\| \leq Ke^{-\kappa(t-s)} \|\epsilon'_s\| \times (1 + \|g\|).$$

Thus the term (27) is dominated by

$$K(1 + \|g\|) \int_0^t e^{-\kappa(t-s)} \|\epsilon'_s\| ds,$$

from which we prove (27) like in the previous lemma. **QED**

4.7 Tightness results

We refer the reader to section 5.1.2 in the appendix, where tightness criteria for families of $C(M)$ -valued random variables are given. They will be used in this section.

4.7.1 Tightness of $(L_{\mu^*}^{-1}(M)_t)_{t \geq 0}$

In this section we prove the following lemma which in particular implies the tightness of $(D_t)_{t \geq 0}$ and of $(D'_t)_{t \geq 0}$.

Lemma 4.12 $(L_{\mu^*}^{-1}(M)_t)_{t \geq 0}$ is tight.

Proof : We have the relation (that defines $L_{\mu^*}^{-1}(M)$)

$$dL_{\mu^*}^{-1}(M)_t(x) = -G_{\mu^*} L_{\mu^*}^{-1}(M)_t(x) dt + dM_t(x).$$

Thus, using the expression of G_{μ^*}

$$dL_{\mu^*}^{-1}(M)_t(x) = -\frac{1}{2} L_{\mu^*}^{-1}(M)_t(x) dt + A_t(x) dt + dM_t(x),$$

with

$$A_t(x) = \widehat{C}_{\mu^*}(V_x, L_{\mu^*}^{-1}(M)_t).$$

Since μ^* is absolutely continuous with respect to λ , we have that

$$\|A_t\| \leq K \|L_{\mu^*}^{-1}(M)_t\|_{\lambda}$$

and therefore (using lemma 4.4 (i) for $\alpha = 2$)

$$\sup_t \mathbb{E}[\|A_t\|^2] < \infty.$$

We also have

$$\text{Lip}(A_t) \leq K \|L_{\mu^*}^{-1}(M)_t\|_{\lambda},$$

where $\text{Lip}(A_t)$ is the Lipschitz constant of A_t (see (38))

In order to prove this tightness result, we first prove that for all x , $(L_{\mu^*}^{-1}(M)_t(x))_t$ is tight. Setting $Z_t^x = L_{\mu^*}^{-1}(M)_t(x)$ we have

$$\begin{aligned} \frac{d}{dt} \mathbb{E}[(Z_t^x)^2] &\leq -\mathbb{E}[(Z_t^x)^2] + 2\mathbb{E}[|Z_t^x| \times |A_t(x)|] + \frac{d}{dt} \mathbb{E}[\langle M(x) \rangle_t] \\ &\leq -\mathbb{E}[(Z_t^x)^2] + K\mathbb{E}[(Z_t^x)^2]^{1/2} + K \end{aligned}$$

which implies that $(L_{\mu^*}^{-1}(M)_t(x))_t$ is bounded in $L^2(\mathbf{P})$ and thus tight.

We now estimate $\mathbb{E}[|Z_t^x - Z_t^y|^\alpha]^{1/\alpha}$ for α greater than 2 and the dimension of M . Setting $Z_t^{x,y} = Z_t^x - Z_t^y$, we have

$$\begin{aligned}
\frac{d}{dt}\mathbb{E}[(Z_t^{x,y})^\alpha] &\leq -\frac{\alpha}{2}\mathbb{E}[(Z_t^{x,y})^\alpha] + \alpha\mathbb{E}[(Z_t^{x,y})^{\alpha-1}|A_t(x) - A_t(y)|] \\
&\quad + \frac{\alpha(\alpha-1)}{2}\mathbb{E}\left[(Z_t^{x,y})^{\alpha-2}\frac{d}{dt}\langle M(x) - M(y)\rangle_t\right] \\
&\leq -\frac{\alpha}{2}\mathbb{E}[(Z_t^{x,y})^\alpha] + \alpha d(x,y)\mathbb{E}[(Z_t^{x,y})^{\alpha-1}\text{Lip}(A_t)] \\
&\quad + Kd(x,y)^2\mathbb{E}[(Z_t^{x,y})^{\alpha-2}] \\
&\leq -\frac{\alpha}{2}\mathbb{E}[(Z_t^{x,y})^\alpha] + Kd(x,y)\mathbb{E}[(Z_t^{x,y})^{\alpha-1}\|L^{-1}(M)_t\|_\lambda] \\
&\quad + Kd(x,y)^2\mathbb{E}[(Z_t^{x,y})^{\alpha-2}] \\
&\leq -\frac{\alpha}{2}\mathbb{E}[(Z_t^{x,y})^\alpha] + Kd(x,y)\mathbb{E}[(Z_t^{x,y})^\alpha]^{\frac{\alpha-1}{\alpha}}\mathbb{E}[\|L^{-1}(M)_t\|_\lambda^\alpha]^{1/\alpha} \\
&\quad + Kd(x,y)^2\mathbb{E}[(Z_t^{x,y})^\alpha]^{\frac{\alpha-2}{\alpha}} \\
&\leq -\frac{\alpha}{2}\mathbb{E}[(Z_t^{x,y})^\alpha] + Kd(x,y)\mathbb{E}[(Z_t^{x,y})^\alpha]^{\frac{\alpha-1}{\alpha}} \\
&\quad + Kd(x,y)^2\mathbb{E}[(Z_t^{x,y})^\alpha]^{\frac{\alpha-2}{\alpha}}.
\end{aligned}$$

Thus, if $x_t = \mathbb{E}[(Z_t^{x,y})^\alpha]/d(x,y)^\alpha$,

$$\frac{dx_t}{dt} \leq -\frac{\alpha}{2}x_t + Kx_t^{\frac{\alpha-1}{\alpha}} + Kx_t^{\frac{\alpha-2}{\alpha}}.$$

It is now an exercise to show that $x_t \leq K$ and so that

$$\mathbb{E}[(Z_t^{x,y})^\alpha]^{1/\alpha} \leq Kd(x,y).$$

Using corollary 5.3, this completes the proof for the tightness of $(L_{\mu^*}^{-1}(M)_t)_t$.

QED

Remark 4.13 *Kolmogorov's theorem (see theorem 1.4.1 and its proof in Kunita (1990)), with the estimates given in the proof of this lemma, implies that*

$$\sup_t \mathbb{E}[\|L_{\mu^*}^{-1}(M)_t\|] < \infty.$$

4.7.2 Tightness of $((L_{\mu^*}^g)^{-1}(N^g, M)_t)_{t \geq 0}$

Fix $g = (g_1, \dots, g_n) \in C(M)^n$. Let $\widehat{\Delta}g$ be defined by the relation

$$(\widehat{\Delta}g, L_{\mu^*}^{-1}(M)) = (L_{\mu^*}^g)^{-1}(N^g, M).$$

Set $A_tg = (A_tg_1, \dots, A_tg_n)$ with $A_tg_i = \widehat{C}_{\mu^*}(g_i, L_{\mu^*}^{-1}(M)_t)$. Then

$$d\widehat{\Delta}_t g = dN_t^g - \frac{\widehat{\Delta}_t g}{2} dt + A_tg dt.$$

Thus,

$$\widehat{\Delta}_t g = e^{-t/2} \int_0^t e^{s/2} dN_s^g + e^{-t/2} \int_0^t e^{s/2} A_s g ds.$$

Using this expression it is easy to prove that $(\widehat{\Delta}_t g)_{t \geq 0}$ is bounded in $L^2(\mathbf{P})$. This implies, using also lemma 4.12

Lemma 4.14 $((L_{\mu^*}^g)^{-1}(N^g, M)_t)_{t \geq 0}$ is tight.

4.8 Convergence in law of $(N^g, M)_{t+} - (N^g, M)_t$

In this section, we denote by $\mathbf{E}_t$ the conditional expectation with respect to $\mathcal{F}_{e^t}$. We also set $\mathbf{Q} = \mathbf{Q}_{\mu^*}$ and $C = \widehat{C}_{\mu^*}$.

4.8.1 Preliminary lemmas.

For $f \in C(M)$ and $t \geq 0$, set $N_s^{f,t} = N_{t+s}^f - N_t^f$.

Lemma 4.15 For all f and g in $C(M)$,

$$\lim_{t \rightarrow \infty} \langle N^{f,t}, N^{g,t} \rangle_s = s \times C(f, g).$$

Proof : Set

$$G(z) = \langle \nabla \mathbf{Q}f, \nabla \mathbf{Q}g \rangle(z) - C(f, g)$$

and

$$G_u(z) = \langle \nabla \mathbf{Q}_u f, \nabla \mathbf{Q}_u g \rangle(z) - C(f, g).$$

We have

$$\begin{aligned}
\langle N^{f,t}, N^{g,t} \rangle_s - s \times C(f, g) &= \int_{e^t}^{e^{t+s}} G_u(X_u) \frac{du}{u} \\
&= \int_{e^t}^{e^{t+s}} (G_u - G)(X_u) \frac{du}{u} \\
&\quad + \int_{e^t}^{e^{t+s}} G(X_u) \frac{du}{u}.
\end{aligned}$$

Integrating by parts, we get that

$$\int_{e^t}^{e^{t+s}} G(X_u) \frac{du}{u} = (\mu_{e^{t+s}} G - \mu_{e^t} G) + \int_0^s (\mu_{e^t+u} G) du.$$

Since $\mu^* G = 0$, this converges towards 0 on the event $\{\mu_t \rightarrow \mu^*\}$. The term $\int_{e^t}^{e^{t+s}} (G_u - G)(X_u) \frac{du}{u}$ converges towards 0 because $(\mu, z) \mapsto \nabla Q_\mu f(z)$ is continuous. This proves the lemma. **QED**

Let $f_1, \dots, f_n$ be in $C(M)$. Let (t_k) be an increasing sequence converging to ∞ such that the conditional law of $M^{n,k} = (N^{f_1, t_k}, \dots, N^{f_n, t_k})$ given $\mathcal{F}_{e^{t_k}}$ converges in law towards a $\mathbb{R}^n$ -valued process $W^n = (W_1, \dots, W_n)$.

Lemma 4.16 *W^n is a centered Gaussian process such that for all i and j ,*

$$\mathbb{E}[W_i^n(s)W_j^n(t)] = (s \wedge t)C(f_i, f_j).$$

Proof : We first prove that W^n is a martingale. For all k , $M^{n,k}$ is a martingale. For all $u \leq v$, Burkholder-Davies-Gundy inequality (BDG inequality in the following) implies that $(M^{n,k}(v) - M^{n,k}(u))_k$ is bounded in L^2 .

Let $l \geq 1$, $\varphi \in C(\mathbb{R}^l)$, $0 \leq s_1 \leq \dots \leq s_l \leq u$ and $(i_1, \dots, i_l) \in \{1, \dots, n\}^l$. Then for all k and $i \in \{1, \dots, n\}$, the martingale property implies that

$$\mathbb{E}_{t_k}[(M_i^{n,k}(v) - M_i^{n,k}(u))Z_k] = 0$$

where Z_k is of the form

$$Z_k = \varphi(M_{i_1}^{n,k}(s_1), \dots, M_{i_l}^{n,k}(s_l)). \tag{28}$$

Using the convergence of the conditional law of $M^{n,k}$ given $\mathcal{F}_{e^{t_k}}$ towards the law of W^n and since $(M_i^{n,k}(v) - M_i^{n,k}(u))_k$ is uniformly integrable (because it is bounded in L^2), we prove that

$$\mathbb{E}[(W_i^n(v) - W_i^n(u))Z] = 0$$

where Z is of the form

$$Z = \varphi(W_{i_1}^n(s_1), \dots, W_{i_l}^n(s_l)). \quad (29)$$

This implies that W^n is a martingale.

We now prove that for $(i, j) \in \{1, \dots, n\}$ (with $C = C_{\mu^*}$),

$$\langle W_i^n, W_j^n \rangle_s = s \times C(f_i, f_j).$$

By definition of $\langle M_i^{n,k}, M_j^{n,k} \rangle$ (in the following $\langle \cdot, \cdot \rangle_u^v = \langle \cdot, \cdot \rangle_v - \langle \cdot, \cdot \rangle_u$)

$$\begin{aligned} \mathbb{E}_{t_k} \left[\left((M_i^{n,k}(v) - M_i^{n,k}(u))(M_j^{n,k}(v) - M_j^{n,k}(u)) \right. \right. \\ \left. \left. - \langle M_i^{n,k}, M_j^{n,k} \rangle_u^v \right) Z_k \right] = 0 \end{aligned} \quad (30)$$

where Z_k is of the form (28). Using the convergence in law and the fact that $(M_i^{n,k}(v) - M_i^{n,k}(u))^2_k$ is bounded in L^2 (still using BDG inequality), we prove that as $k \rightarrow \infty$,

$$\mathbb{E}_{t_k} [(M_i^{n,k}(v) - M_i^{n,k}(u))(M_j^{n,k}(v) - M_j^{n,k}(u))Z_k]$$

converges towards

$$\mathbb{E}[(W_i^n(v) - W_i^n(u))(W_j^n(v) - W_j^n(u))Z].$$

with Z of the form (29). Now,

$$\begin{aligned} \mathbb{E}_{t_k} [\langle M_i^{n,k}, M_j^{n,k} \rangle_v Z_k] - v \times \mathbb{E}[Z] \times C(x_i, x_j) \\ = \mathbb{E}_{t_k} [(\langle M_i^{n,k}, M_j^{n,k} \rangle_v - v \times C(f_i, f_j))Z_k] \\ + v \times (\mathbb{E}_{t_k}[Z_k] - \mathbb{E}[Z]) \times C(f_i, f_j) \end{aligned}$$

The convergence in L^2 of $\langle M_i^{n,k}, M_j^{n,k} \rangle_v$ towards $v \times C(f_i, f_j)$ shows that the first term converges towards 0. The convergence of the conditional law of $M^{n,k}$ with respect to $\mathcal{F}_{e^{t_k}}$ towards W^n shows that the second term converges towards 0. Thus

$$\mathbb{E} [((W_i^n(v) - W_i^n(u))(W_j^n(v) - W_j^n(u)) - (v - u)C(f_i, f_j)) Z] = 0.$$

This shows that $\langle W_i^n, W_j^n \rangle_s = s \times C(f_i, f_j)$. We conclude using Lévy's theorem. **QED**

4.8.2 Convergence in law of $M_{t+} - M_t$

In this section, we denote by $\mathcal{L}_t$ the conditional law of $M_{t+} - M_t$ knowing $\mathcal{F}_{e^t}$. Then $\mathcal{L}_t$ is a probability measure on $C(\mathbb{R}^+ \times M)$.

Proposition 4.17 *When $t \rightarrow \infty$, $\mathcal{L}_t$ converges weakly towards the law of a $C(M)$ -valued Brownian motion of covariance C_{μ^*} .*

Proof : In the following, we will simply denote $M_{t+} - M_t$ by M^t . We first prove that

Lemma 4.18 *$\{\mathcal{L}_t : t \geq 0\}$ is tight.*

Proof : For all $x \in M$, t and u in $\mathbb{R}^+$,

$$\begin{aligned} \mathbb{E}_t[(M_u^t(x))^2] &= \mathbb{E}_t \left[\int_t^{t+u} d\langle M(x) \rangle_s \right] \\ &\leq Ku. \end{aligned}$$

This implies that for all $u \in \mathbb{R}^+$ and $x \in M$, $(M_u^t(x))_{t \geq 0}$ is tight.

Let $\alpha > 0$. We fix $T > 0$. Then for (u, x) and (v, y) in $[0, T] \times M$, using BDG inequality,

$$\begin{aligned} \mathbb{E}_t[|M_u^t(x) - M_v^t(y)|^\alpha]^\frac{1}{\alpha} &\leq \mathbb{E}_t[|M_u^t(x) - M_u^t(y)|^\alpha]^\frac{1}{\alpha} \\ &\quad + \mathbb{E}_t[|M_u^t(y) - M_v^t(y)|^\alpha]^\frac{1}{\alpha} \\ &\leq K_\alpha \times (\sqrt{T}d(x, y) + \sqrt{|v - u|}) \end{aligned}$$

where K_α is a positive constant depending only on α , $\|V\|$ and $\text{Lip}(V)$ the Lipschitz constant of V .

We now let D_T be the distance on $[0, T] \times M$ defined by

$$D_T((u, x), (v, y)) = K_\alpha \times (\sqrt{T}d(x, y) + \sqrt{|v - u|}).$$

The covering number $N([0, T] \times M, D_T, \epsilon)$ is of order $\epsilon^{-d-1/2}$ as $\epsilon \rightarrow 0$. Taking $\alpha > d + 1/2$, we conclude using corollary 5.3. **QED**

Let (t_k) be an increasing sequence converging to ∞ and N a $C(M)$ -valued random process (or a $C(\mathbb{R}^+ \times M)$ random variable) such that $\mathcal{L}_{t_k}$ converges in law towards N .

Lemma 4.19 *N is a $C(M)$ -valued Brownian motion of covariance C_{μ^*} .*

Proof : Let W be a $C(M)$ -valued Brownian motion of covariance C_{μ^*} . Using lemma 4.16, we prove that for all $(x_1, \dots, x_n) \in M^n$, $(N(x_1), \dots, N(x_n))$ has the same distribution as $(W(x_1), \dots, W(x_n))$. This implies the lemma. **QED**

Since $\{\mathcal{L}_t\}$ is tight, this lemma implies that $\mathcal{L}_t$ converges weakly towards the law of a $C(M)$ -valued Brownian motion of covariance C_{μ^*} . **QED**

4.8.3 Convergence in law of $(N^g, M)_{t+} - (N^g, M)_t$

In this section, we fix $g = (g_1, \dots, g_n) \in C(M)^n$ and we denote by $\mathcal{L}_t^g$ the conditional law of $(N^g, M)_{t+} - (N^g, M)_t$ knowing $\mathcal{F}_{et}$. Then $\mathcal{L}_t^g$ is a probability measure on $C(\mathbb{R}^+ \times M \cup \{1, \dots, n\})$. In the following we will denote $(N^{g,t}, M^t)$ the process $(N^g, M)_{t+} - (N^g, M)_t$.

Let $(W_t^f)_{(t,f) \in \mathbb{R}^+ \times C(M)}$ be a $\mathcal{X}(M)$ -valued Brownian motion of covariance $\widehat{C}_{\mu^*}$. Denoting $W_t(x) = W_t^{V_x}$, then $W = (W_t(x))_{(t,x) \in \mathbb{R}^+ \times M}$ is a $C(M)$ -valued Brownian motion of covariance C_{μ^*} . For $g = (g_1, \dots, g_n) \in C(M)^n$, W^g will denote $(W^{g_1}, \dots, W^{g_n})$. In the following we will simply denote (W^g, W) the process $(W_t^g, (W_t(x))_{x \in M})_{t \geq 0}$.

Proposition 4.20 *As t goes to ∞ , $\mathcal{L}_t^g$ converges weakly towards the law of (W^g, W) .*

Proof : We first prove that

Lemma 4.21 *$\{\mathcal{L}_t^g : t \geq 0\}$ is tight.*

Proof : This is a straightforward consequence of the tightness of $\{\mathcal{L}_t\}$ and of the fact that for all $\alpha > 0$, there exists K_α such that for all nonnegative u and v , $\mathbb{E}_t[|N_u^{g,t} - N_v^{g,t}|^\alpha]^\frac{1}{\alpha} \leq K_\alpha \sqrt{|v - u|}$. **QED**

Let (t_k) be an increasing sequence converging to ∞ and $(\tilde{N}^g, \tilde{M})$ a $\mathbb{R}^n \times C(M)$ -valued random process (or a $C(\mathbb{R}^+ \times M \cup \{1, \dots, n\})$ random variable) such that $\mathcal{L}_{t_k}^g$ converges in law towards $(\tilde{N}^g, \tilde{M})$. Then lemmas 4.15 and 4.16 imply that $(\tilde{N}^g, \tilde{M})$ has the same law as (W^g, W) . Since $\{\mathcal{L}_t^g\}$ is tight, $\mathcal{L}_t^g$ converges towards the law of (W^g, W) . **QED**

4.9 Convergence in law of D

4.9.1 Convergence in law of $(D_{t+s} - e^{-sG_{\mu^*}} D_t)_{s \geq 0}$

We have

$$D'_{t+s} - e^{-sG_{\mu^*}} D'_t = L_{\mu^*}^{-1}(M^t)_s + \int_0^s e^{-(s-u)G_{\mu^*}} \epsilon'_{t+u} du.$$

Since (using lemma 4.9)

$$\left\| \int_0^s e^{-(s-u)G_{\mu^*}} \epsilon'_{t+u} du \right\| \leq K e^{-\kappa t}$$

and $\|D_t - D'_t\| \leq K(1+t)e^{-t/2}$, this proves that $(D_{t+s} - e^{-sG_{\mu^*}} D_t - L_{\mu^*}^{-1}(M_{t+} - M_t)_s)_{s \geq 0}$ converges towards 0. Since $L_{\mu^*}^{-1}$ is continuous, this proves that the law of $L_{\mu^*}^{-1}(M_{t+} - M_t)$ converges weakly towards $L_{\mu^*}^{-1}(W)$. Since $L_{\mu^*}^{-1}(W)$ is an Ornstein-Uhlenbeck process of covariance C_{μ^*} and drift $-G_{\mu^*}$ started from 0, we have

Theorem 4.22 *The conditional law of $(D_{t+s} - e^{-sG_{\mu^*}} D_t)_{s \geq 0}$ given $\mathcal{F}_{e^t}$ converges weakly towards an Ornstein-Uhlenbeck process of covariance C_{μ^*} and drift $-G_{\mu^*}$ started from 0.*

4.9.2 Convergence in law of D_{t+}

We can now prove theorem 3.1. We here denote by P_t the semigroup of an Ornstein-Uhlenbeck process of covariance C_{μ^*} and drift $-G_{\mu^*}$, and we denote by π its invariant probability measure.

We know that (as $t \rightarrow \infty$) $(D_{t+s} - e^{-sG_{\mu^*}} D_t)_{s \geq 0}$ converges in law towards $L_{\mu^*}^{-1}(W)$, where W is a $C(M)$ -valued Brownian motion of covariance C_{μ^*} . Since $(D_t)_{t \geq 0}$ is tight, there exists $\nu \in \mathcal{P}(C(M))$ and an increasing sequence t_n converging towards ∞ such that D_{t_n} converges in law towards ν . Then D_{t_n+} converges in law towards $(L_{\mu^*}^{-1}(W)_s + e^{-sG_{\mu^*}} Z_0)$, with Z_0 independent of W and distributed like ν . This proves that D_{t_n+} converges in law towards an Ornstein-Uhlenbeck process of covariance C_{μ^*} and drift $-G_{\mu^*}$.

We now fix $t > 0$. Let s_n be a subsequence of t_n such that D_{s_n-t+} converges in law. Then D_{s_n-t} converges towards a law we denote by ν_t and D_{s_n-t+} converges in law towards an Ornstein-Uhlenbeck process of covariance C_{μ^*} and drift $-G_{\mu^*}$. Since $D_{s_n} = D_{s_n-t+t}$, D_{s_n} converges in law towards $\nu_t P_t$. On the other hand D_{s_n} converges in law towards ν . Thus $\nu_t P_t = \nu$.

Let φ be a Lipschitz bounded function on $C(M)$. Then

$$\begin{aligned} |\nu_t \mathbf{P}_t \varphi - \pi \varphi| &= \left| \int (\mathbf{P}_t \varphi(f) - \pi \varphi) \nu_t(df) \right| \\ &\leq \int |\mathbf{P}_t \varphi(f) - \mathbf{P}_t \varphi(0)| \nu_t(df) + |\mathbf{P}_t \varphi(0) - \pi \varphi| \end{aligned}$$

where the second term converges towards 0 (using (37)) and the first term is dominated by (using lemma 5.15)

$$K e^{-\kappa t} \int \|f\| \nu_t(df).$$

It is easy to check that

$$\begin{aligned} \int \|f\| \nu_t(df) &= \lim_{k \rightarrow \infty} \int (\|f\| \wedge k) \nu_t(df) \\ &= \lim_{k \rightarrow \infty} \lim_{n \rightarrow \infty} \mathbf{E}[\|D_{s_n-t}\| \wedge k] \\ &\leq \lim_{n \rightarrow \infty} \mathbf{E}[\|D_{s_n-t}\|] \\ &\leq \sup_t \mathbf{E}[\|D_t\|]. \end{aligned}$$

Since

$$\begin{aligned} \|D_t\| &\leq \|D_t^1 + D_t^5\| + \|L_{\mu^*}^{-1}(M)_t\| \\ &\quad + \left\| \int_0^t e^{(t-s)G_{\mu^*}} \epsilon'_s ds \right\|, \end{aligned}$$

using the estimates (22), the proof of lemma 4.10 and remark 4.13, we get that

$$\sup_{t \geq 0} \mathbf{E}[\|D_t\|] < \infty.$$

Taking the limit, we prove $\nu \varphi = \pi \varphi$ for all Lipschitz bounded function φ on $C(M)$. This implies $\nu = \pi$, which proves the theorem. **QED**

4.9.3 Convergence in law of D^g

We can also prove theorem 3.2.

For $g = (g_1, \dots, g_n) \in C(M)^n$, we set $D_t^g = (\Delta_t g, D_t)$, and $D_t'^g = (\Delta_t' g, D_t')$. Since $\|D_t^g - D_t'^g\| \leq K(1+t)e^{-t/2}$, instead of studying D^g , we can only study $D_t'^g$. Then

$$\begin{aligned} D_{t+s}'^g - e^{-sG_{\mu^*}^g} D_t'^g &= (L_{\mu^*}^g)^{-1} (N^{g,t}, M^t)_s \\ &\quad + \int_0^s e^{-(s-u)G_{\mu^*}^g} (\epsilon'_{t+u} g, \epsilon'_{t+u}) du. \end{aligned}$$

The norm of the second term of the right hand side (using the proof of lemma 4.11) is dominated by

$$\begin{aligned} &\leq K(1 + \|g\|) \int_0^s e^{-\kappa(s-u)} \|\epsilon'_{t+u}\| du \\ &\leq K \int_0^s e^{-\kappa(s-u)} (1+t+u)^2 e^{-(t+u)/2} du \\ &\leq k e^{-\kappa t} \end{aligned}$$

Like in section 4.9.1, since $(L_{\mu^*}^g)^{-1} (W^g, W)$ is an Ornstein-Uhlenbeck process of covariance $C_{\mu^*}^g$ and drift $-G_{\mu^*}^g$ started from 0,

Theorem 4.23 *The conditional law of $((\Delta^g, D)_{t+s} - e^{-sG_{\mu^*}^g} (\Delta^g, D)_t)_{s \geq 0}$ given $\mathcal{F}_{e^t}$ converges weakly towards an Ornstein-Uhlenbeck process of covariance $C_{\mu^*}^g$ and drift $-G_{\mu^*}^g$ started from 0.*

From this theorem, like in section 4.9.2, we prove theorem 3.2. **QED**

5 Appendix : Random variables and Ornstein-Uhlenbeck processes on $C(M)$

5.1 $C(M)$ -valued random variables

5.1.1 Generalities

Let (M, d) be a compact metric space (note that there is no assumption here that M is a manifold), $C(M)$ the space of real valued continuous functions on M equipped with the uniform norm $\|f\| = \sup_{x \in M} |f(x)|$. By classical results, $C(M)$ is a separable (see e.g. [16]) Banach space (see e.g. [11] or [9]) and its topological dual is the space $\mathcal{M}(M)$ of bounded signed measures on

M (see e.g [11] or [9]). For $\mu \in \mathcal{M}(M)$ and $f \in C(M)$ we use the notation $\mu f = \langle \mu, f \rangle = \int_M f d\mu$.

Let $(\Omega, \mathcal{F}, \mathbf{P})$ be a probability space. A $C(M)$ -valued random variable is a Borel map $F : \Omega \rightarrow C(M)$.

For $x \in M$, let $\pi_x : C(M) \rightarrow \mathbb{R}$, denote the projection defined by

$$\pi_x(f) = f(x).$$

Lemma 5.1 *The Borel σ -field on $C(M)$ is the σ -field generated by the maps $\{\pi_x\}_{x \in M}$. In particular*

- (i) *A map $F : \Omega \rightarrow C(M)$ is a $C(M)$ -valued random variable if and only if $\{\pi_x(F)\}_{x \in M}$ is a family of real valued random variables.*
- (ii) *The law of a $C(M)$ -valued random variable is determined by its finite dimensional distributions (i.e the law of $\{\pi_x(F)\}_{x \in I}$ with $I \subset M$ finite).*

Proof : Let $\mathcal{A} = \sigma\{\pi_x, x \in M\}$ and $\mathcal{B}$ the Borel σ -field on $C(M)$. The maps π_x being continuous, $\mathcal{B}$ contains $\mathcal{A}$. Conversely, let $B_f(r) = \{g \in C(M) : \|g - f\| \leq r\}$ and let S be a countable dense subset of M . Then $B_f(r) = \cap_{x \in S} \{g \in C(M) : |\pi_x(f) - \pi_x(g)| \leq r\}$ Hence $B_f(r) \in \mathcal{A}$. Since $C(M)$ is separable, $\mathcal{B}$ is generated by the sets $\{B_f(r), f \in C(M), r \geq 0\}$. **QED**

5.1.2 Tightness criteria

Let $\mathcal{P}(C(M))$ be the space of Borel probability measures on $C(M)$. An element ν of $\mathcal{P}(C(M))$ is the law of a $C(M)$ -valued random variable F , and $\nu = \mathbf{P}_F$. Recall that a sequence $\{\nu^n\}$ in $\mathcal{P}(C(M))$ is said *converging weakly* towards $\nu \in \mathcal{P}(C(M))$ if $\int \varphi d\nu_n \rightarrow \int \varphi d\nu$ for every bounded and continuous function $\varphi : C(M) \rightarrow \mathbb{R}$. A sequence $\{F_n\}$ of $C(M)$ -valued random variable is said *converging in law* towards F a $C(M)$ -valued random variable if $\{\mathbf{P}_{F_n}\}$ converges in law towards $\mathbf{P}_F$. A family $\mathcal{X} \subset \mathcal{P}(C(M))$ is said to be *tight* if for every $\epsilon > 0$ there exists some compact set $\mathcal{K} \subset C(M)$ such that $\mathbf{P}(\mathcal{K}) \geq 1 - \epsilon$ for all $\mathcal{P} \in \mathcal{X}$. A family of random variables is said to be tight if the family of their laws is tight.

Since $C(M)$ is a separable and complete, Prohorov theorem [8] asserts that $\mathcal{X} \subset \mathcal{P}(C(M))$ is tight if and only if it is relatively compact.

The next proposition gives a useful criterium for a class of random variables to be tight. It follows directly from [14] (Corollary 11.7 p. 307 and the remark following Theorem 11.2). A function $\psi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is a Young function if it is convex, increasing and $\psi(0) = 0$. If Z is a real valued random variable, we let

$$\|Z\|_\psi = \inf\{c > 0 : \mathbf{E}(\psi(|Z|/c)) \leq 1\}.$$

For $\epsilon > 0$, we denote by $N(M, d; \epsilon)$ the covering number of E by balls of radius less than ϵ (i.e. the minimal number of balls of radius less than ϵ that cover E), and by D the diameter of M .

Proposition 5.2 *Let $(F_t)_{t \in I}$ be a family of $C(M)$ -valued random variables and ψ a Young function. Assume that*

- (i) *There exists $x \in E$ such that $(F_t(x))_{t \in I}$ is tight;*
- (ii) $\|F_t(x) - F_t(y)\|_\psi \leq Kd(x, y)$;
- (iii) $\int_0^D \psi^{-1}(N(M, d; \epsilon))d\epsilon < \infty$.

Then $(F_t)_{t \geq 0}$ is tight.

Corollary 5.3 *Suppose M is a compact finite dimensional manifold of dimension r , d the Riemannian distance, and*

$$[\mathbf{E}|F_t(x) - F_t(y)|^\alpha]^{1/\alpha} \leq Kd(x, y)$$

for some $\alpha > r$. Then conditions (ii) and (iii) of Proposition 5.2 hold true.

Proof : One has $N(E, d; \epsilon)$ is of order ϵ^{-r} ; and for $\psi(x) = x^\alpha$, $\|\cdot\|_\psi$ is the L^α norm. Hence the result. **QED**

5.1.3 $C(M)$ -valued Gaussian variable

Recall that a (centered) real-valued random variable Y with variance σ^2 is said to be *Gaussian* if it has distribution

$$P_Y(dx) = \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{x^2}{2\sigma^2}\right)dx.$$

Its characteristic function is then

$$\Phi_Y(t) = \mathbb{E}[\exp(itY)] = \exp\left(-\frac{t^2\sigma^2}{2}\right).$$

Here we adopt the convention that the zero function ($Y = 0$) is Gaussian with variance 0 and that all the Gaussian random variables are centered.

A family $\{Y_i\}_{i \in I}$ of real-valued random variables is said to be Gaussian if for all finite set $J \subset I$ and for all $\alpha \in \mathbb{R}^J$, $\sum_{j \in J} \alpha_j Y_j$ is Gaussian.

A $C(M)$ -valued random variable F is said to be *Gaussian* if for all $\mu \in \mathcal{M}(M)$, $\langle \mu, F \rangle$ is Gaussian.

Lemma 5.4 *A $C(M)$ -valued random variable F is Gaussian if and only if the family $\{\pi_x(F)\}$ is Gaussian.*

Proof : The direct implication is obvious. We prove the second. Assume that $\{\pi_x(F)\}$ is a Gaussian family. Let μ be a probability over M . By the strong law of large number and the separability of $C(M)$ there exists a nonempty set $\Lambda \subset M^{\mathbb{N}}$ (actually Λ has $\mu^{\mathbb{N}}$ measure 1) such that for all $(x_i) \in \Lambda$ and all $f \in C(M)$

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n f(x_i) = \langle \mu, f \rangle.$$

In particular $Y_n \rightarrow \langle \mu, F \rangle$ where

$$Y_n = \frac{1}{n} \sum_{i=1}^n F(x_i).$$

And, by Lebesgue theorem, $\Phi_{Y_n}(t) \rightarrow \Phi_{\langle \mu, F \rangle}(t)$. Since, by assumption Y_n is Gaussian, $\Phi_{Y_n}(t) = \exp(-t^2 \sigma_n^2 / 2)$. Let $\sigma \in [0, \infty]$ be a limit point of (σ_n) . Then $\Phi_{\langle \mu, F \rangle}(t) = \exp(-t^2 \sigma^2 / 2)$. This proves that $\sigma < \infty$ (a characteristic function being continuous) and that $\langle \mu, F \rangle$ is Gaussian.

If now $\mu \in \mathcal{M}(M)$ by Jordan-Hann decomposition we may write $\mu = a\mu_1 - b\mu_2$ with $a, b \geq 0$ and μ_1, μ_2 probabilities. It follows from what precede that $\langle \mu, X \rangle$ is Gaussian. **QED**

Given a $C(M)$ -valued Gaussian random variable F we let $\mathbf{Var}_F : \mathcal{M}(M) \rightarrow \mathbb{R}$ denote the *variance function* of F defined by

$$\mathbf{Var}_F(\mu) = \mathbf{E}(\langle \mu, F \rangle^2).$$

In view of lemma 5.1 (ii), the law of F is entirely determined by its variance function.

A useful property of Gaussian variables is the following.

Lemma 5.5 *Let M' be another compact metric space and $A : C(M) \rightarrow C(M')$ a bounded linear operator. Let F be a $C(M)$ -valued Gaussian random variable. Then AF is a $C(M')$ -valued Gaussian random variable with variance*

$$\mathbf{Var}_{AF} = \mathbf{Var}_F \circ A^*$$

where $A^* : \mathcal{M}(M') \rightarrow \mathcal{M}(M)$ is the adjoint of A .

Proof : follows from the duality $\langle \mu, AF \rangle = \langle A^*\mu, F \rangle$ and the definitions. **QED**

5.2 Brownian motions on $C(M)$.

Let $C : M \times M \rightarrow \mathbb{R}$ be a continuous symmetric (i.e $C(x, y) = C(y, x)$) function such that

$$\sum_{ij} a_i a_j C(x_i, x_j) \geq 0$$

for every finite sequence (a_i, x_i) with $a_i \in \mathbb{R}$ and $x_i \in M$. Such a function is sometimes called a *Mercer kernel*.

A *Brownian motion on $C(M)$ with covariance C* is a $C(M)$ -valued stochastic process $W = \{W_t\}_{t \geq 0}$ such that $W_0 = 0$ and for each $T \geq 0$, $W^T = \{W_t(x) : t \leq T, x \in M\}$ is a $C([0, T] \times M)$ -valued Gaussian random variable with variance

$$\mathbf{Var}_{W^T}(\nu) = \int_{([0, T] \times M)^2} (s \wedge t) C(x, y) \nu(ds dx) \nu(dt dy)$$

or equivalently

$$\mathbb{E}(W_t(x)W_s(y)) = (s \wedge t)C(x, y).$$

Let

$$d_C(x, y) = \sqrt{C(x, x) - 2C(x, y) + C(y, y)}.$$

The d_C is a pseudo-distance on M . For $\epsilon > 0$, let

$$\omega_C(\epsilon) = \sup\{\eta > 0 : d(x, y) \leq \eta \Rightarrow d_C(x, y) \leq \epsilon\}.$$

Then $N(M, d; \omega_C(\epsilon)) \geq N(M, d_C; \epsilon)$.

Hypothesis 5.6

$$\int_0^1 \log(N(d, M; \omega_C(\epsilon))) d\epsilon < \infty$$

where $N(d, M; \eta)$ is the covering number of M by balls of radius less than η .

Remark 5.7 Assume that M is a compact finite dimensional manifold and that $d_C(x, y) \leq Kd(x, y)^\alpha$ for some $\alpha > 0$. Then $\omega_C(\epsilon) \leq (\frac{\epsilon}{K})^{1/\alpha}$ and $N(d, M; \eta) = O(\eta^{-\dim(M)})$; so that the preceding hypothesis holds.

Proposition 5.8 Under hypothesis 5.6 there exists a Brownian motion on $C(M)$ with covariance C .

Proof : By Mercer Theorem (see e.g. [10]) there exists a countable family of function $\Psi_i \in C(M), i \in \mathbb{N}$, such that

$$C(x, y) = \sum_i \Psi_i(x)\Psi_i(y)$$

and the convergence is uniform. Let $B^i, i \in \mathbb{N}$, be a family of independent standard Brownian motions. Set

$$W_t^n(x) = \sum_{i \leq n} B_t^i \Psi_i(x), n \geq 0.$$

Then, for each $(t, x) \in \mathbb{R}^+ \times M$, the sequence $(W_t^n(x))_{n \geq 1}$ is a Martingale. It is furthermore bounded in L^2 since

$$\mathbb{E}[(W_t^n(x))^2] = t \sum_{i \leq n} \Psi_i(x)^2 \leq tC(x, x).$$

Hence by Doob's convergence theorem one may define

$$W_t(x) = \sum_{i \geq 0} B_t^i \Psi_i(x).$$

Let now $S \subset \mathbb{R}^+ \times M$ be a countable and dense set. It is easily checked that the family $(W_t(x))_{(t,x) \in S}$ is a centered Gaussian family with covariance given by

$$\mathbb{E}[W_s(x)W_t(y)] = (s \wedge t)C(x, y),$$

In particular, for $t \geq s$

$$\begin{aligned} \mathbb{E}[(W_s(x) - W_t(y))^2] &= sC(x, x) - 2sC(x, y) + tC(y, y) \\ &= sd_C(x, y)^2 + (t - s)C(y, y) \\ &\leq K(t - s) + sd_C(x, y)^2 \end{aligned}$$

This later bound combined with classical results on Gaussian processes (see e.g Theorem 11.17 in [14]) implies that $(t, x) \mapsto W_t(x)$ admits a version uniformly continuous over $S_T = \{(t, x) \in S : t \leq T\}$. By density it can be extended to a continuous (in (t, x)) process

$$W = (W_t(x))_{\{(t,x) \in \mathbb{R}^+ \times M\}}$$

The process W can be viewed as a $C(M)$ -valued continuous random process with the desired covariance. **QED**

5.3 Ornstein-Uhlenbeck processes

Let $A : C(M) \rightarrow C(M)$ be a bounded operator and W a $C(M)$ -valued Brownian motion with covariance C as defined in the preceding section.

An *Ornstein-Uhlenbeck process* with drift A , covariance C and initial condition $F_0 = f \in C(M)$ is defined to be a $C(M)$ valued stochastic process continuous in t , such that

$$F_t - f = \int_0^t AF_s ds + W_t. \tag{31}$$

Note that we may think of F as the solution to the “stochastic differential equation” on $C(M)$:

$$dF_t = AF_t dt + dW_t$$

with initial condition $F_0 = f \in C(M)$.

Our aim here is to construct such a solution and state some of its properties.

We let $(e^{tA})_{t \in \mathbb{R}}$ denote the linear flow induced by A . Recall that for each t , e^{tA} is the bounded operator on $C(M)$ defined by

$$e^{tA} = \sum_{k \in \mathbb{N}} \frac{(tA)^k}{k!}.$$

Given $T > 0$ we let $L_A : C(\mathbb{R}^+ \times M) \rightarrow C(\mathbb{R}^+ \times M)$ be defined by

$$L_A(f)_t = f_t - f_0 - \int_0^t A f_s ds, \quad t \geq 0. \quad (32)$$

Given $T > 0$ we let $L_A^T : C([0, T] \times M) \rightarrow C([0, T] \times M)$ be defined by

$$L_A^T(f)_t = f_t - f_0 - \int_0^t A f_s ds, \quad 0 \leq t \leq T. \quad (33)$$

Note that if for $f \in C(\mathbb{R}^+ \times M)$, we let $f^T \in C([0, T] \times M)$ be defined by $f_t^T = f_t$ for $t \in [0, T]$.

Lemma 5.9 L_A^T is a bounded operator and its restriction to $C_0([0, T] \times M) = \{f \in C([0, T] \times M) : f_0 = 0\}$ is bijective with inverse $(L_A^T)^{-1}$ defined by

$$(L_A^T)^{-1}(g)_t = g_t + \int_0^t e^{(t-s)A} A g_s ds, \quad t \in [0, T]. \quad (34)$$

Proof : Linearity of L_A^T is obvious. Also

$$\|L_A^T(f)\| \leq (2 + T\|A\|)\|f\|.$$

This proves that L_A^T is bounded.

Observe that $L_A^T(f) = 0$ implies that $f_t = e^{tA} f_0$. Hence L_A^T restricted to $C_0([0, T] \times M)$ is injective. Let $g \in C_0([0, T] \times M)$ and let f_t be given by the left hand side of (34). Then

$$h_t = L_A^T(f)_t - g_t = \int_0^t e^{(t-s)A} A g_s ds - \int_0^t A f_s ds.$$

It is easily seen that h is differentiable and that $\frac{d}{dt} h_t = 0$. This proves that $h_t = h_0 = 0$. **QED**

We also have

Lemma 5.10 *The restriction of L_A to $C_0(\mathbb{R}^+ \times M) = \{f \in C(\mathbb{R}^+ \times M) : f_0 = 0\}$ is bijective with inverse $(L_A)^{-1}$ defined by*

$$L_A^{-1}(g)_t = g_t + \int_0^t e^{(t-s)A} A g_s ds. \quad (35)$$

The next lemma easily follows.

Lemma 5.11 *For all $f \in C(M)$ and $g \in C_0(\mathbb{R}^+ \times M)$ the solution to*

$$f_t = f + \int_0^t A f_s ds + g_t,$$

is given by

$$f_t = e^{tA} f + L_A^{-1}(g)_t.$$

If now W is a $C(M)$ -valued Brownian motion as defined in the preceding section, one may define

$$F_t = e^{tA} f + L_A^{-1}(W)_t.$$

Such a process is the unique solution to (31). Note that, by Lemma 5.5 $(F_t - e^{tA} f)_{t \leq T}$ is a $C_0([0, T] \times M)$ -valued Gaussian random variable. In particular

Proposition 5.12 *Let (F_t) be the solution to (31) with initial condition $F_0 = 0$. Then for each $t \geq 0$, F_t is a $C(M)$ -valued Gaussian random variable with variance*

$$\text{Var}_{F_t}(\mu) = \int_0^t \langle \mu, e^{sA} C e^{sA*} \mu \rangle ds.$$

where $C : \mathcal{M}(M) \rightarrow C(M)$ is the operator defined by $C\mu(x) = \int_M C(x, y)\mu(dy)$.

Proof : Fix $T > 0$. To shorten notation let $G : C_0([0, T] \times M) \rightarrow C(M)$ be the operator defined by

$$G(g) = (L_A^T)^{-1}(g)_T.$$

Hence $F_T = G(W^T)$. By Lemma 5.5, F_T is Gaussian with variance

$$\text{Var}_{F_T} = \text{Var}_{W^T} \circ G^*.$$

Now, for all $\nu \in \mathcal{M}([0, T] \times M)$

$$\text{Var}_{W^T}(\nu) = \langle \nu, \mathcal{C}\nu \rangle$$

where

$$\mathcal{C}\nu(s, x) = \int_{[0, T] \times M} (s \wedge u) C(x, y) \nu(dudy).$$

Thus

$$\text{Var}_{F_T}(\mu) = \langle \mu, G\mathcal{C}G^*\mu \rangle.$$

Our next goal is to compute $G\mathcal{C}G^*\mu$. It easily follows from the definition of G that

$$G^*\mu = \delta_T \otimes \mu + dsA^*e^{(T-s)A^*}\mu.$$

Thus (integrating by parts)

$$\mathcal{C}G^*\mu = \mathcal{C}\nu_1 + \mathcal{C}\nu_2$$

with $\nu_1 = \delta_T \otimes \mu$ and $\nu_2 = dsA^*e^{(T-s)A^*}\mu$. One has

$$\mathcal{C}\nu_1(s, x) = s(C\mu)(x);$$

$$\mathcal{C}\nu_2(s, x) = \int_0^T (s \wedge u)(C\dot{m}_u(\mu))(x)du$$

with $m_u(\mu) = -e^{(T-u)A^*}\mu$ and $\dot{m}_u(\mu)$ stands for the derivative of $u \mapsto m_u(\mu)$.

Thus

$$\mathcal{C}\nu_2(s, x) = -sC\mu - \int_0^s Cm_u(\mu)(x)du$$

and

$$\mathcal{C}G^*\mu(s, x) = -\int_0^s Cm_u(\mu)(x)du.$$

Set

$$h_s(x) = \int_0^s Cm_u(\mu)(x)du.$$

Then

$$\begin{aligned} G\mathcal{C}G^*\mu &= h_T + \int_0^T e^{(T-s)A}Ah_sds = \int_0^T e^{(T-s)A}\dot{h}_sds \\ &= \int_0^T e^{(T-s)A}Ce^{(T-s)A^*}\mu ds = \int_0^T e^{sA}Ce^{sA^*}\mu ds. \end{aligned}$$

QED

5.3.1 Asymptotic Behaviour

Let $\lambda(A) = \lim_{t \rightarrow \infty} \frac{\log(\|e^{tA}\|)}{t}$ which exists by subadditivity. Then for some constant $K < \infty$, $\|e^{tA}\| \leq Ke^{\lambda(A)t}$ for all positive t . Let (F_t) denote the solution to (31), with $F_0 = f \in C(M)$.

Corollary 5.13 *Assume $\lambda(A) < 0$. Then for each $\mu \in \mathcal{M}(M)$ $(\langle \mu, F_t \rangle)$ converges in law toward a Gaussian random variable with variance*

$$V(\mu) = \int_0^\infty \langle \mu, e^{sA} C e^{sA*} \mu \rangle ds.$$

Proof : follows from proposition 5.12 and Lemma 5.11 **QED**

Corollary 5.14 *Assume that $\lambda(A) < 0$. Set*

$$d_V(x, y) = \sqrt{V(\delta_x - \delta_y)}$$

and

$$\omega_V(\epsilon) = \sup\{\eta > 0 : d(x, y) \leq \eta \Rightarrow d_V(x, y) \leq \epsilon\}.$$

Assume furthermore that ω_V verifies the condition expressed by hypothesis 5.6. Then (F_t) converges in law toward a $C(M)$ -valued Gaussian random variable with variance V .

Proof : Let ν_t denote the law of F_t . Corollary 5.13 and lemma 5.1 imply that every limit point of $\{\nu_t\}$ (for the weak* topology) is the law of a $C(M)$ -valued Gaussian variable with variance V . The proof then reduces to show that (ν_t) is relatively compact or equivalently that $\{F_t\}$ is tight. We use Proposition 5.2. The first condition follows from Lemma 5.13. Let $\psi(x) = e^{x^2} - 1$. It is easily verified that for any real valued Gaussian random variable Z with variance σ^2 , $\|Z\|_\Psi = \sigma\sqrt{8/3}$. Hence $\|F_t(x) - F_t(y)\|_\psi \leq 2d_V(x, y)$ so that condition (ii) holds with the pseudo distance d_V . By definition of ω_V , $N(M, d; \omega_V(\epsilon)) \geq N(M, d_V; \epsilon)$ and since $\psi^{-1}(u) = \sqrt{\log(u-1)}$ condition (iii) is verified. **QED**

Denote by P_t the semigroup associated to an Ornstein-Uhlenbeck process of covariance C and drift A . Then for all bounded measurable $\varphi : C(M) \rightarrow \mathbb{R}$ and $f \in C(M)$,

$$P_t \varphi(f) = E[\varphi(F_t)]. \tag{36}$$

Denote by π the law of a $C(M)$ -valued Gaussian random variable with variance V . Then π is the invariant probability measure of P_t , i.e. $\pi P_t = \pi$. Corollary 5.14 implies that, when $\lambda(A) < 0$, for all $f \in C(M)$ and all bounded continuous $\varphi : C(M) \rightarrow \mathbb{R}$,

$$\lim_{t \rightarrow \infty} P_t \varphi(f) = \pi \varphi. \quad (37)$$

Even though we don't have the speed of convergence in the previous limit, we have

Lemma 5.15 *Assume that $\lambda(A) < 0$. For all bounded Lipschitz continuous $\varphi : C(M) \rightarrow \mathbb{R}$, all f and g in $C(M)$,*

$$|P_t \varphi(f) - P_t \varphi(g)| \leq K e^{\lambda(A)t} \|f - g\|.$$

Proof : We have $P_t \varphi(f) = E[\varphi(L_A^{-1}(W)_t + e^{tA}f)]$. So, using the fact that φ is Lipschitz,

$$\begin{aligned} |P_t \varphi(f) - P_t \varphi(g)| &\leq K \|e^{tA}(f - g)\| \\ &\leq K e^{\lambda(A)t} \|f - g\|. \end{aligned}$$

This proves the lemma. **QED**

To conclude this section we give a set of simple sufficient conditions ensuring that the hypotheses of corollary 5.14 are satisfied.

For $f \in C(M)$ we let

$$\text{Lip}(f) = \sup_{x \neq y} \frac{|f(x) - f(y)|}{d(x, y)} \in \mathbb{R}^+ \cup \{\infty\}. \quad (38)$$

A map f is said to be Lipschitz provided $\text{Lip}(f) < \infty$.

Proposition 5.16 *Assume*

- (i) $N(d, M; \epsilon) = O(\epsilon^{-r})$ for some $r > 0$ (This holds in particular if M is a finite dimensional manifold).
- (ii) $x \mapsto C(x, y)$ is Lipschitz uniformly in y . That is

$$\sup_{z \in M} |C(x, z) - C(y, z)| \leq K d(x, y)$$

for some $K \geq 0$.

(iii) *There exists $K > 0$ such that*

$$Lip(Af) \leq K(Lip(f) + \|f\|).$$

(iv) $\lambda(A) < 0$

Then the hypotheses, hence the conclusion, of corollary 5.14 are satisfied.

We begin with the following lemma.

Lemma 5.17 *Under hypotheses (iii) and (iv) of proposition 5.16*

$$Lip(e^{tA}f) \leq e^{Kt}(Lip(f) + K'\|f\|)$$

for some constants K, K' .

Proof : For all x, y

$$\begin{aligned} |e^{tA}f(x) - e^{tA}f(y)| &= \left| \int_0^t [Ae^{sA}f(x) - Ae^{sA}f(y)]ds + f(x) - f(y) \right| \\ &\leq K \left(\int_0^t [Lip(e^{sA}f) + \|e^{sA}f\|] ds + Lip(f) \right) d(x, y). \end{aligned}$$

Since $\lambda = \lambda(A) < 0$, there exists $C > 0$ such that $\|e^{sA}\| \leq Ce^{-s\lambda}$. Thus

$$Lip(e^{tA}f) \leq K \int_0^t Lip(e^{sA}f)ds + \frac{KC}{\lambda}\|f\| + Lip(f)$$

and the result follows from Gronwal's lemma. **QED**

We now pass to the proof of the proposition. In what follows the constants may change from line to line.

Proof : Set $\mu = \delta_x - \delta_y$ and $f_s = Ce^{sA^*}\mu$ so that

$$\langle \mu, e^{sA}Ce^{sA^*}\mu \rangle = e^{sA}f_s(x) - e^{sA}f_s(y).$$

It follows from hypotheses (ii) and (iv) that

$$Lip(f_s) + \|f_s\| \leq Ke^{-s\lambda}$$

for some positive constants K and a . Therefore, by the preceding lemma,

$$Lip(e^{sA}f_s) \leq Ke^{s\alpha}$$

for some (other) positive constants K, α . Thus

$$\begin{aligned}
d_V(x, y)^2 &\leq d(x, y) \int_0^T \text{Lip}(e^{sA} f_s) ds + \int_T^\infty (e^{sA} f(x) - e^{sA} f(y)) ds \\
&\leq d(x, y) \int_0^T K e^{s\alpha} ds + 2 \int_T^\infty \|e^{sA} f_s\| ds \\
&\leq K \left(d(x, y) e^{\alpha T} + \int_T^\infty e^{-s\lambda} ds \right) \\
&\leq K (d(x, y) e^{\alpha T} + e^{-\lambda T}).
\end{aligned}$$

Let $\gamma = \frac{\alpha}{\lambda}$, $\epsilon > 0$, and $T = -\ln(\epsilon)/\lambda$. Then

$$d_V^2(x, y) \leq K(\epsilon^{-\gamma} d(x, y) + \epsilon).$$

Therefore

$$d(x, y) \leq \epsilon^{\gamma+1} \Rightarrow d_V^2(x, y) \leq K\epsilon,$$

so that $N(d, M; \omega_V(\epsilon)) = O(\epsilon^{-2r(\gamma+1)})$ and hypothesis 5.6 holds true. **QED**

Example 5.18 Let

$$Af(x) = \int f(y) k(x, dy)$$

with

$$k(x, dy) = k_0(x, y) \mu(dy) + \sum_{i=1}^n a_i(x) \delta_{b_i(x)}$$

where

- (i) μ is a bounded measure on M ,
- (ii) $k_0(x, y)$ is bounded and uniformly Lipschitz in x ,
- (iii) $a_i : M \rightarrow \mathbb{R}$ and $b_i : M \rightarrow M$ are Lipschitz.

Then hypothesis (iii) of proposition 5.16 is verified.

References

- [1] J.R. Baxter, G.A. Brosamler, (1976), *Energy and the law of iterated logarithm*, Math. Scand. **38**, 115–136.
- [2] J.R. Baxter, G.A. Brosamler, (1983), *Recurrence of Brownian motions on compact manifolds*. In Colloquium in honor of Laurent Schwartz, Vol. 2 (Palaiseau, 1983). Astérisque No. 132 (1985), 15–46.
- [3] M. Benaïm, M. Ledoux and O. Raimond, (2002), *Self-interacting diffusions*, Probab. Theor. Relat. Fields **122**, 1-41.
- [4] M. Benaïm and O. Raimond, (2003), *Self-interacting diffusions II: Convergence in Law.*, Annales de l’institut Henri-Poincaré **6**, 1043-1055.
- [5] M. Benaïm and O. Raimond, (2005), *Self-interacting diffusions III: Symmetric interactions.*, Annals of Probability **33**, no. 5, 1717–1759.
- [6] M. Benaïm and O. Raimond, (2009), *A Bakry-Emery Criterion for self-interacting diffusions.*, Seminar on Stochastic Analysis, Random Fields and Applications V, 19–22, Progr. Probab., 59, Birkhauser, Basel.
- [7] R.N. Bhattacharya, (1982) *On the functional central limit theorem and the law of the iterated logarithm for Markov processes*. Z. Wahrsch. Verw. Gebiete, **60**(2):185-201.
- [8] P. Billingsley, (1968) “Convergence of Probability Measures”, Wiley Series in Probability and Mathematical Statistics, John Wiley and Sons.
- [9] J.B. Conway, (1985), “A course in Functional Analysis”, Graduate texts in Mathematics, Springer Verlag.
- [10] J. Dieudonné, (1972) “Eléments d’Analyse I,” Gautier-Villars, Paris.
- [11] R. Dudley, (2003), “Real Analysis and Probability” Cambridge studies in advanced mathematics, Cambridge University Press.
- [12] M. Duflo, (1996), “Algorithmes Stochastiques”, Mathématiques et Applications, Springer-Verlag, vol 23.
- [13] H. Kunita, (1990), ”Stochastic Flows and Stochastic Differential Equations”, Cambridge Univ. Press.

- [14] M. Ledoux and M. Talagrand, (1991) "Probability in Banach spaces", Springer (1991). Second printing (2002).
- [15] R. Pemantle, (2007), *A survey of random processes with reinforcement*, Probability survey, Vol 4, 1-79.
- [16] D. Stroock, (1993), "Probability, an Analytic view", Cambridge Univ. Press.